

**CHARACTERIZATION OF RIPARIAN MANAGEMENT ZONES AND
UPLAND MANAGEMENT AREAS WITH RESPECT
TO WILDLIFE HABITAT**

1989 FIELD REPORT

By

Washington Department of Wildlife
Habitat Management Division




October 1990



**Washington Department of Wildlife
Habitat Management Division
Timber-Fish-Wildlife Project
TFW-003-90-003**

1989 FIELD REPORT



**Characterization of Riparian
Management Zones and Upland
Management Areas with Respect
to Wildlife Habitat**

October 1990

1989 FIELD REPORT

CHARACTERIZATION OF
RIPARIAN MANAGEMENT ZONES
&
UPLAND MANAGEMENT AREAS
WITH RESPECT TO WILDLIFE HABITAT

Submitted to:

Washington Department of Natural Resources
Division of Forest Regulation and Assistance
1007 S. Washington St., Mail Stop EL-03
Olympia, WA 98504

Submitted By:

TFW Wildlife Steering Committee
under the direction of the
Cooperative Monitoring, Evaluation, and Research Committee

Prepared by:

Andy Carlson
TFW Biologist
Washington Department of Wildlife
Habitat Management Division
600 Capitol Way N., Mail Stop GJ-11
Olympia, WA 98501-1091
October 23, 1990

This report summarizes the 1988 and 1989 field seasons of the Cooperative, Monitoring, Evaluation, and Research Committee research project #3 titled: "Characterization of Riparian Management Zones and Upland Management Areas with Respect to Wildlife Habitat". In December of 1990 it was decided by the Wildlife Steering Committee that a final report would not be produced for the 1989 field season. Instead of producing a final report a summary of the data collected is presented in this 1989 Field Report. The Wildlife Steering Committee has given their approval of the 1989 Field Report with limited editing.

Planning is currently taking place to produce a cumulative report summarizing data collected from 1988 to 1990. The 1988-90 cumulative report will be available in the Spring of 1991.

The opinions, findings, conclusions, or recommendations expressed in this report are those of the authors and do not necessarily reflect the views of any participant in, or committee of, the Timber/Fish/Wildlife Agreement, the Washington Forest Practices Board, or the Washington Department of Natural Resources, nor does mention of trade names or commercial products constitute endorsement or recommendation for use.

TABLE OF CONTENTS

LIST OF TABLES	iii
LIST OF FIGURES	xi
ABSTRACT	1
INTRODUCTION	2
STUDY AREA	3
METHODS	3
SITE SELECTION	3
DATA ANALYSIS	4
RMZ/UMA SITE SUMMARY	6
RMZ RESULTS	12
LARGE ORGANIC DEBRIS	12
VEGETATION AND OTHER STRIP VARIABLES	14
LIVE TREE DENSITY	60
SNAG DENSITY	67
UMA RESULTS	73
VEGETATION AND OTHER STRIP VARIABLES	73
LIVE TREE DENSITY	96
SNAG DENSITY	100
RECOMMENDATIONS	103
ACKNOWLEDGEMENTS	104
LITERATURE CITED	105

APPENDIX A 107

List of species codes, scientific,
and common names of trees,
shrubs, and herbs.

APPENDIX B 114

Key contacts: Source for forest
practice information.

LIST OF TABLES

Table LOD-1. Eastside boulder/bedrock RMZ average number of large organic debris pieces per 100 feet.	12
Table LOD-2. Westside boulder/bedrock RMZ average number of large organic debris pieces per 100 feet.	13
Table LOD-3. Eastside gravel/cobble RMZ average number of large organic debris pieces per 100 feet.	13
Table LOD-4. Westside gravel/cobble RMZ average number of large organic debris pieces per 100 feet.	13
Table SHRUB-1. Eastside lake RMZs, water type 1, dominant shrub #1 mean subplot coverage and constancy.	14
Table SHRUB-2. Eastside lake RMZs, water type 2, dominant shrub #1 mean subplot coverage and constancy.	15
Table SHRUB-3. Eastside lake RMZs, water type 1, dominant shrub #2 mean subplot coverage and constancy.	15
Table SHRUB-4. Westside lake RMZs, water type 1, dominant shrub #1 mean subplot coverage and constancy.	16
Table SHRUB-5. Westside lake RMZs, water type 2, dominant shrub #1 mean subplot coverage and constancy.	17
Table SHRUB-6. Westside lake RMZs, water type 3, dominant shrub #1 mean subplot coverage and constancy.	17
Table SHRUB-7. Westside lake RMZs, water type 1, dominant shrub #2 mean subplot coverage and constancy.	18
Table SHRUB-8. Westside lake RMZs, water type 2, dominant shrub #2 mean subplot coverage and constancy.	19
Table SHRUB-9. Westside lake RMZs, water type 3, dominant shrub #2 mean subplot coverage and constancy.	19
Table SHRUB-10. Eastside, boulder/bedrock, water type 3, dominant shrub #1 mean subplot coverage and constancy.	20
Table SHRUB-11. Westside, boulder/bedrock, water type 1, dominant shrub #1 mean subplot coverage and constancy.	21
Table SHRUB-12. Westside, boulder/bedrock, water type 2, dominant shrub #1 mean subplot coverage and constancy.	22

Table SHRUB-13. Westside, boulder/bedrock, water type 3, dominant shrub #1 mean subplot coverage and constancy.	22
Table SHRUB-14. Westside, boulder/bedrock, water type 1, dominant shrub #2 mean subplot coverage and constancy.	23
Table SHRUB-15. Westside, boulder/bedrock, water type 2, dominant shrub #2 mean subplot coverage and constancy.	24
Table SHRUB-16. Westside, boulder/bedrock, water type 3, dominant shrub #2 mean subplot coverage and constancy.	24
Table SHRUB-17. Eastside, gravel/cobble, water type 1, dominant shrub #1 mean subplot coverage and constancy.	25
Table SHRUB-18. Eastside, gravel/cobble, water type 2, dominant shrub #1 mean subplot coverage and constancy.	25
Table SHRUB-19. Eastside, gravel/cobble, water type 3, dominant shrub #1 mean subplot coverage and constancy.	26
Table SHRUB-20. Eastside, gravel/cobble, water type 1, dominant shrub #2 mean subplot coverage and constancy.	27
Table SHRUB-21. Eastside, gravel/cobble, water type 3, dominant shrub #2 mean subplot coverage and constancy.	28
Table SHRUB-22. Westside, gravel/cobble, water type 1, dominant shrub #1 mean subplot coverage and constancy.	29
Table SHRUB-23. Westside, gravel/cobble, water type 2, dominant shrub #1 mean subplot coverage and constancy.	30
Table SHRUB-24. Westside, gravel/cobble, water type 3, dominant shrub #1 mean subplot coverage and constancy.	31
Table SHRUB-25. Westside, gravel/cobble, water type 1, dominant shrub #2 mean subplot coverage and constancy.	32
Table SHRUB-26. Westside, gravel/cobble, water type 2, dominant shrub #2 mean subplot coverage and constancy.	33
Table SHRUB-27. Westside, gravel/cobble, water type 3, dominant shrub #2 mean subplot coverage and constancy.	34
Table HERB-1. Eastside lake RMZs, water type 1, dominant herb #1 mean subplot coverage and constancy.	35
Table HERB-2. Eastside lake RMZs, water type 1, dominant herb #2 mean subplot coverage and constancy.	36

Table HERB-3. Westside lake RMZs, water type 1, dominant herb #1 mean subplot coverage and constancy.	37
Table HERB-4. Westside lake RMZs, water type 2, dominant herb #1 mean subplot coverage and constancy.	37
Table HERB-5. Westside lake RMZs, water type 3, dominant herb #1 mean subplot coverage and constancy.	38
Table HERB-6. Westside lake RMZs, water type 1, dominant herb #2 mean subplot coverage and constancy.	39
Table HERB-7. Westside lake RMZs, water type 2, dominant herb #2 mean subplot coverage and constancy.	40
Table HERB-8. Westside lake RMZs, water type 3, dominant herb #2 mean subplot coverage and constancy.	40
Table HERB-9. Westside, boulder/bedrock, water type 1, dominant herb #1 mean subplot coverage and constancy.	41
Table HERB-10. Westside, boulder/bedrock, water type 2, dominant herb #1 mean subplot coverage and constancy.	41
Table HERB-11. Westside, boulder/bedrock, water type 3, dominant herb #1 mean subplot coverage and constancy.	42
Table HERB-12. Westside, boulder/bedrock, water type 1, dominant herb #2 mean subplot coverage and constancy.	43
Table HERB-13. Westside, boulder/bedrock, water type 2, dominant herb #2 mean subplot coverage and constancy.	43
Table HERB-14. Westside, boulder/bedrock, water type 3, dominant herb #2 mean subplot coverage and constancy.	44
Table HERB-15. Eastside, gravel/cobble, water type 1, dominant herb #1 mean subplot coverage and constancy.	45
Table HERB-16. Eastside, gravel/cobble, water type 3, dominant shrub #1 mean subplot coverage and constancy.	45
Table HERB-17. Eastside, gravel/cobble, water type 1, dominant herb #2 mean subplot coverage and constancy.	46
Table HERB-18. Eastside, gravel/cobble, water type 3, dominant shrub #2 mean subplot coverage and constancy.	47
Table HERB-19. Westside, gravel/cobble, water type 1, dominant herb #1 mean subplot coverage and constancy.	48
Table HERB-20. Westside, gravel/cobble, water type 2, dominant herb #1 mean subplot coverage and constancy.	49

Table HERB-21. Westside, gravel/cobble, water type 3, dominant herb #1 mean subplot coverage and constancy.	50
Table HERB-22. Westside, gravel/cobble, water type 1, dominant herb #2 mean subplot coverage and constancy.	51
Table HERB-23. Westside, gravel/cobble, water type 2, dominant herb #2 mean subplot coverage and constancy.	52
Table HERB-24. Westside, gravel/cobble, water type 3, dominant herb #2 mean subplot coverage and constancy.	53
Table COVER-1. Eastside lake RMZ mean coverage/constancy for subplot canopy, total shrubs, forbs, and graminoids.	54
Table COVER-2. Westside lake RMZ mean coverage/constancy for subplot canopy, total shrubs, forbs, and graminoids.	55
Table COVER-3. Eastside boulder/bedrock RMZ mean coverage/constancy for subplot canopy, total shrubs, forbs, and graminoids.	55
Table COVER-4. Westside boulder/bedrock RMZ mean coverage/constancy for subplot canopy, total shrubs, forbs, and graminoids.	56
Table COVER-5. Eastside gravel/cobble RMZ mean coverage/constancy for subplot canopy, total shrubs, forbs, and graminoids.	56
Table COVER-6. Westside gravel/cobble RMZ mean coverage/constancy for subplot canopy, total shrubs, forbs, and graminoids.	57
Table TREE-1. Eastside lake RMZ mean tree density all water types - conifers.	61
Table TREE-2. Eastside lake RMZ mean tree density all water types - hardwoods.	61
Table TREE-3. Westside lake RMZ mean tree density all water types - conifers.	62
Table TREE-4. Westside lake RMZ mean tree density all water types - hardwoods.	62
Table TREE-5. Eastside boulder/bedrock RMZ mean tree density all water types - conifers.	63
Table TREE-6. Eastside boulder/bedrock RMZ mean tree density all water types - hardwoods.	63

Table TREE-7. Westside boulder/bedrock RMZ mean tree density all water types - conifers.	64
Table TREE-8. Westside boulder/bedrock RMZ mean tree density all water types - hardwoods.	64
Table TREE-9. Eastside gravel/cobble RMZ mean tree density all water types - conifers.	65
Table TREE-10. Eastside gravel/cobble RMZ mean tree density all water types - hardwoods.	65
Table TREE-11. Westside gravel/cobble RMZ mean tree density all water types - conifers.	66
Table TREE-12. Westside gravel/cobble RMZ mean tree density all water types - hardwoods.	66
Table SNAG-1. Eastside lake RMZ mean snag density all water types - conifers.	67
Table SNAG-2. Eastside lake RMZ mean snag density all water types - hardwoods.	67
Table SNAG-3. Westside lake RMZ mean snag density all water types - conifers.	68
Table SNAG-4. Westside lake RMZ mean snag density all water types - hardwoods.	68
Table SNAG-5. Eastside boulder/bedrock RMZ mean snag density all water types - conifers.	69
Table SNAG-6. Eastside boulder/bedrock RMZ mean snag density all water types - hardwoods.	69
Table SNAG-7. Westside boulder/bedrock RMZ mean snag density all water types - conifers.	69
Table SNAG-8. Westside boulder/bedrock RMZ mean snag density all water types - hardwoods.	70
Table SNAG-9. Eastside gravel/cobble RMZ mean snag density all water types - conifers.	70
Table SNAG-10. Eastside gravel/cobble RMZ mean snag density all water types - hardwoods.	71
Table SNAG-11. Westside gravel/cobble RMZ mean snag density all water types - conifers.	71
Table SNAG-12. Westside gravel/cobble RMZ mean snag density all water types - hardwoods.	72

Table SHRUB-28. Eastside UMAs, forested wetlands, dominant shrub #1 mean subplot coverage and constancy.	74
Table SHRUB-29. Eastside UMAs, upland forest, dominant shrub #1 mean subplot coverage and constancy.	75
Table SHRUB-30. Eastside UMAs, forested wetlands, sub-dominant shrub #1 mean subplot coverage and constancy.	76
Table SHRUB-31. Eastside UMAs, upland forest, sub-dominant shrub #1 mean subplot coverage and constancy.	77
Table SHRUB-32. Westside UMA, bogs, dominant shrub #1 mean subplot coverage and constancy.	78
Table SHRUB-33. Westside UMAs, forested wetlands, dominant shrub #1 mean subplot coverage and constancy.	79
Table SHRUB-34. Westside UMAs, upland forest, dominant shrub #1 mean subplot coverage and constancy.	80
Table SHRUB-35. Westside UMA, bogs, dominant shrub #2 mean subplot coverage and constancy.	81
Table SHRUB-36. Westside UMAs, forested wetlands, dominant shrub #2 mean subplot coverage and constancy.	82
Table SHRUB-37. Westside UMAs, upland forest, dominant shrub #2 mean subplot coverage and constancy.	83
Table HERB-25. Eastside UMAs, forested wetlands, dominant herb #1 mean subplot coverage and constancy.	84
Table HERB-26. Eastside UMAs, upland forest, dominant herb #1 mean subplot coverage and constancy.	85
Table HERB-27. Eastside UMAs, forested wetlands, dominant herb #2 mean subplot coverage and constancy.	86
Table HERB-28. Eastside UMAs, upland forest, dominant herb #2 mean subplot coverage and constancy.	87
Table HERB-29. Westside UMA, bogs, dominant herb #1 mean subplot coverage and constancy.	88
Table HERB-30. Westside UMAs, forested wetlands, dominant herb #1 mean subplot coverage and constancy.	89
Table HERB-31. Westside UMAs, upland forest, dominant herb #1 mean subplot coverage and constancy.	89
Table HERB-32. Westside UMA, bogs, dominant herb #2 mean subplot coverage and constancy.	90

Table HERB-33. Westside UMAs, forested wetlands, dominant herb #2 mean subplot coverage and constancy.	92
Table HERB-34. Westside UMAs, upland forest, dominant herb #2 mean subplot coverage and constancy.	93
Table UMACOVER-1. Eastside UMA mean coverage/constancy for subplot canopy, total shrubs, forbs, and graminoids.	94
Table UMACOVER-2. Westside UMA mean coverage/constancy for subplot canopy, total shrubs, forbs, and graminoids.	94
Table UMA-1. Eastside, forested wetland, UMA mean tree density - conifers.	96
Table UMA-2. Eastside, forested wetland, UMA mean tree density - hardwoods.	97
Table UMA-3. Eastside, upland forest, UMA mean tree density - conifers.	97
Table UMA-4. Eastside, upland forest, UMA mean tree density - hardwoods.	97
Table UMA-5. Westside, forested wetland, UMA mean tree density - conifers.	98
Table UMA-6. Westside, forested wetland, UMA mean tree density - hardwoods.	98
Table UMA-7. Westside, upland forest, UMA mean tree density - conifers.	98
Table UMA-8. Westside, upland forest, UMA mean tree density - hardwoods.	98
Table UMA-9. Westside, bog, UMA mean tree density - conifers.	99
Table UMA-10. Westside, bog, UMA mean tree density - hardwoods.	99
Table UMA-11. Eastside, forested wetland, UMA mean snag density - conifers.	100
Table UMA-12. Eastside, forested wetland, UMA mean snag density - hardwoods.	100
Table UMA-13. Eastside, upland forest, UMA mean snag density - conifers.	100
Table UMA-14. Eastside, upland forest, UMA mean snag density - hardwoods.	101

Table UMA-15. Westside, forested wetland, UMA mean snag density - conifers.	101
Table UMA-16. Westside, forested wetland, UMA mean snag density - hardwoods.	101
Table UMA-17. Westside, upland forest, UMA mean snag density - conifers.	102
Table UMA-18. Westside, upland forest, UMA mean snag density - hardwoods.	102
Table UMA-19. Westside, bog, UMA mean snag density - conifers.	102
Table UMA-20. Westside, bog, UMA mean snag density - hardwoods.	102

LIST OF FIGURES

Figure 1. Location of sample sites.	6
Figure 2. Total RMZ and UMA sites sampled.	7
Figure 3. Total RMZ and UMA acres sampled.	8
Figure 4. Site ownerships.	9
Figure 5. Average site width by water type.	10
Figure 6. Number of sites sampled by category.	11

ABSTRACT

In June of 1988 the Washington Department of Wildlife (WDW) entered into a research agreement with the the Washington Department of Natural Resources (DNR) in which WDW agreed to inventory Riparian Management Zones (RMZs) and Upland Management Areas (UMAs) throughout the state of Washington. The intent of the Wildlife Steering Committee when designing this project was to provide detailed information on RMZs and UMAs, but not to identify statistical or casual relationships. The objective was to quantify the physical and botanical characteristics of RMZs and UMAs with respect to wildlife habitat. This report summarizes the first (1988) and second (1989) years of a six-year study on state and private commercial forests in Washington. Three hundred and fifty-nine acres of RMZs located on 114 sites were sampled in 1988 and 1989. A total of 80 RMZs were located on industrial forestland, 21 on private non-industrial land, and 13 on state land. One hundred and twenty-six acres of UMAs located on 30 sites were sampled in 1988 and 1989. A total of 26 UMAs were located on industrial forest land, 2 on private non-industrial, and 2 on state lands. The UMAs sampled are a structurally diverse array of forest types ranging from wetlands to old-growth forests. Tabular reports presented were derived from data collected during the 1988 and 1989 field seasons. The 1988 field season lasted three months (Aug. - Oct.). The 1989 field season lasted six months (May - Oct.). Recommendations to improve sampling efficiency and accuracy are provided at the end of this report.

INTRODUCTION

The Timber/Fish/Wildlife (TFW) Agreement (1987) requires the development of a monitoring, evaluation, and research program with cooperative decisions on priorities and associated costs. Results from research and monitoring will be used to make incremental changes in the forest practices regulations. This process is known as adaptive management and is a policy of the Forest Practices Board.

This project (Cooperative Monitoring, Evaluation, and Research Committee Project #3) was designed to provide detailed information on RMZs and UMAs. It is not designed to identify statistical or causal relationships between habitat and wildlife, nor does it attempt to measure compliance with the Forest Practices Act. It provides information for determining effectiveness of the TFW process in protecting riparian zones. The project quantifies the physical and botanical characteristics of RMZs and UMAs with respect to wildlife habitat.

Mean RMZ width and UMA acreages were derived from methods described in WDW's Field Procedures Handbook (Second Edition, 1990).

RMZs are defined in the Forest Practice Regulations, WAC 222 (1988) as a specified area alongside Type 1, 2, and 3 waters where specific measures are taken to protect water quality and fish and wildlife habitat. Riparian zones are among the most heavily used wildlife habitats in the forests of Washington (Thomas et al., 1979). They occur along rivers, streams, intermittent drainages, ponds, lakes, reservoirs, springs, and wetlands.

UMAs are areas of naturally occurring trees and vegetation or where specific silvicultural activities have been designed for wildlife management (Forest Practices Board Manual, 1988). UMAs are voluntary under the TFW agreement. They are intended to accommodate site-specific needs of landowners and wildlife. UMAs are intended to increase wildlife habitat diversity by providing conditions that would not normally occur in timber-harvested areas, such as shelter, corridors for travel, and security for other wildlife activities associated with harvest areas. The TFW intent was that UMAs would provide increased diversity through irregular scattering or dispersion of habitats for a broad spectrum of wildlife species.

This project provides an information base for more detailed studies on the value and use of RMZs and UMAs for wildlife. The Department of Ecology (Ed Rashin, 206-586-5291) in Olympia is currently conducting a study to monitor the effect RMZs have on water temperature regulation. Department of Ecology study sites are limited to Project #3's sample sites .

This is the second year of a six-year study.

STUDY AREA

This study was limited to commercial state and private forests of Washington. Most western Washington forests are located in the Sitka spruce (*Picea sitchensis*) and western hemlock (*Tsuga heterophylla*) zones. East of the Cascade crest the forests are located in the Douglas-fir (*Pseudotsuga menziesii*), Pacific silver fir (*Abies amabilis*), and subalpine fir (*Abies lasiocarpa*) zones. Franklin and Dyrness (1973) have published an excellent description of the physiography, geology, soils, and climate of this region.

METHODS

The Field Procedures Handbook Second Edition (WDW, 1990) outlines the sampling procedures used to quantify RMZs and UMAs.

Mean RMZ width and UMA acreages were derived from methods described in WDW's Field Procedures Handbook (Second Edition, 1990).

SITE SELECTION

Because sites were often selected as they became available, true stratified random sampling was not possible. To reduce bias in the site selection the following procedure was used:

Sites sampled were limited to harvested areas meeting the requirements of the TFW Agreement of February 1988. Sites meeting TFW standards, but which were harvested prior to February of 1988, were also sampled. The intent was to provide an unbiased, stratified, view of RMZs/UMAs as they occurred throughout the state of Washington. RMZs sampled were limited to those that occur on type 1, 2, and 3 waters.

Water types are defined as follows:

Type 1 waters are those waters inventoried as "shorelines of the state" under chapter 90.58 RCW. Type 2 waters are those waters diverted for domestic use by more than 100 persons, used by substantial numbers of anadromous or resident game fish for spawning, rearing or migration with a defined channel of more than 20 feet, and a gradient of less than four percent. Type 3 waters are those waters diverted for domestic use by more than 10 persons, used by substantial numbers of anadromous or resident game fish for spawning, rearing or migration with a defined channel of

more than five feet, a gradient of less than 12 percent, and are highly significant for protection of downstream water quality.

The Department of Revenue maintains a list of Forest Practices Applications (FPAs) on which timber tax has been paid. FPAs from this list were then collected from individual DNR Regional Offices. These FPAs were screened to select those which contain either RMZs or UMAs.

Concurrently, FPAs containing RMZs/UMAs were also requested from private land-owners (industrial and non-industrial), and Washington Department of Wildlife regional biologists. Using these other sources allowed sampling of RMZs and UMAs that may not have been listed on original FPAs.

FPAs were mapped statewide to display RMZ and UMA locations. From this map, a sampling schedule was established. Emphasis was placed on sampling new areas, according to the annual schedule shown below, as required by contract.

Subsequent years' samples will include a mix of new and older RMZs and UMAs as follows:

Year 1 - (1988) 39 new areas sampled

Year 2 - (1989) 105 new areas sampled

Year 3 - new areas and 20% of 1st year areas

Year 4 - new areas and 20% of 2nd year areas

Year 5 - new areas, 20% of 1st year areas, and 20% of 3rd year areas

Year 6 - new areas, 20% of 2nd year areas, and 20% of 4th year areas

DATA ANALYSIS

Data were originally compiled in a SMARTWARE database (Informix Software, Version 3.1). They were then transferred to PARADOX (Borland, Version 3.0). All tabular summaries were created with Quattro Pro (Borland, Version 1.0). Graphics displayed in the Final Report were produced with Harvard Graphics (Software Publishing Corp., Version 2.12). The final report was produced in Ventura Publisher (Xerox, Version 2.0).

Data summaries were created by the following categorical break downs within the state: Eastern WA or Western WA as defined by the Washington Forest Practices Rules and Regulations (1988), water type (or UMA type) and substrate.

All sample site locations were recorded on 7.5-minute USGS quadrant maps. Sites were recorded on 15-minute maps when 7.5- minute maps were unavailable. A stereo pair of aerial photographs have been filed together with the original field forms, harvest unit maps, and the forest practice application. Maps and files are stored at the Department of Wildlife, Habitat Management Division, 600 Capitol Way N., Olympia, Washington, 98501-1091, (206) 753-3318.

All discussions within this report pertain to sites sampled during the 1988 and 1989 field seasons. Summaries provided are of data collected by Project #3.

RMZ/UMA SITE SUMMARY

Figure 1 maps sample site locations for the 1988/89 field seasons. During the 1988/89 field seasons 114 RMZs and 30 UMAs were sampled (Figure 2). The total acreage of RMZs sampled equaled 359 and the total acreage of UMAs equaled 126 (Figure 3).




Figure 1. Map of RMZ and UMA sample sites.




Figure 2. Total RMZ and UMA sites sampled 1988 and 1989.




Figure 3. Total RMZ and UMA acres 1988 and 1989.

The majority of sample sites were located on private industrial land followed by private non-industrial, and state owned land (Figure 4).



Figure 4. Total sites by owner code.

RMZ average widths are listed in Figure 5. These results are the mean widths of RMZs based on the project's criteria for measuring the physical and botanical characteristics of these sites. These averages should not be used for checking compliance with forest practices regulations.




Figure 5. Average site width by water type 1988 and 1989.

Eastside average RMZ widths are estimated to average 30 feet due to the difficulty of defining RMZ boundaries within partial cut harvest units. When harvest boundaries were not easily identified the sampling effort concluded at 30 feet.

Figure 6. Break down of the total number of sites sampled within each category.

<u>Lakes</u>			
<u>Water Type</u>	<u>1</u>	<u>2</u>	<u>3</u>
Eastside	1	1	
Westside	5	3	1
<u>Boulder/Bedrock</u>			
Eastside			1
Westside	10	2	4
<u>Gravel/Cobble</u>			
Eastside	1	1	11
Westside	11	12	50

<u>Total Number of UMAs Within Each UMA Type By Side</u>			
<u>UMA Type</u>	<u>Forested Wetland</u>	<u>Bog</u>	<u>Upland Forest</u>
Eastside	1		2
Westside	7	2	18

RESULTS

RMZs

Three hundred and fifty-nine acres of RMZs located on 114 sites were sampled in 1988/89.

RMZs were broken down into 12 categories (for data analysis and display) in the following manner: first by their location within the state (eastside or westside), secondly by their water type (1,2,3), and lastly by the stream bed substrate (gravel/cobble or boulder/bedrock). On about five sites the entire RMZ identified on the Forest Practice Application was not sampled due to time limitations.

To be classified as a gravel/cobble substrate 50% of the dominant stones must be less than 10 inches in diameter. The substrate is classified as boulder/bedrock when more than 50% of the dominant stones are greater than 10 inches in diameter.

RMZ summaries are provided in the following order: Average number of large organic debris pieces per 100 feet, dominant shrub mean coverage and constancies, dominant herb mean coverage and constancies, mean coverage and constancy values for overstory canopy closure, total shrubs, forbs, and graminoids, live tree density, and lastly snag densities.

LARGE ORGANIC DEBRIS (LOD)

Table LOD-1. Eastside Boulder/Bedrock RMZ
Average Number of Large Organic Debris Pieces
Per Hundred Feet (Note: only water type 3 RMZs
have been sampled within this category).

WATER TYPE	1	2	3
Average Number of LOD pieces/100 Feet	N.A.	N.A.	4
Number of Sites	N.A.	N.A.	1

Table LOD-2. Westside Boulder/Bedrock RMZ
Average Number of Large Organic Debris Pieces
Per Hundred Feet.

WATER TYPE	1	2	3
Average Number of LOD pieces/100 Feet	4	3	4
Number of Sites	10	1	4

Table LOD-3. Eastside Gravel/Cobble RMZ
Average Number of Large Organic Debris Pieces
Per Hundred Feet.

Water Type	1	2	3
Average Number of LOD pieces/100 Feet	1	2	4
Number of Sites	1	1	11

Table LOD-4. Westside Gravel/Cobble RMZ
Average Number of Large Organic Debris Pieces
Per Hundred Feet.

Water type	1	2	3
Average Number of LOD pieces/100 Feet	4	7	6
Number of Sites	11	9	50

Westside gravel/cobble streams appeared to contain more pieces of LOD per 100 feet than similar eastside streams. Only one eastside boulder/bedrock stream was sampled (water type 3). This stream contained the same average pieces of LOD per hundred feet as westside type 3 streams. On both sides of the state, and within both substrate types, LOD was more frequently found in type 3 streams. LOD was least frequently found within type 1 streams.

VEGETATION AND OTHER STRIP VARIABLES

Data were collected on the two dominant shrubs and herbs, total shrubs, forbs and graminoids (grass), downed wood 1 to 3 (decay class 1 = recent fallen, decay class 3 = rotten), water, rock, and soil. Mean coverage and constancy values were calculated for these variables.

Canopy is defined as the percent of closed canopy above the sample plot. Coverage is defined as the percentage of ground, when viewed from above the subplot, the variable covers within the sample plot. Sample plots are 5x10 feet. Constancy is defined as the degree of presence a variable has within sample plots. Subplot coverage and constancy values are given in percent.

RMZ shrubs and herbs are listed in order by their constancy values. Shrub tables 1 through 27 and herb tables 1 through 24 list the 20 most frequently encountered shrubs or forbs. When fewer than 20 shrubs or forbs are listed, this implies that fewer than 20 were encountered within that specific category. **Values are given in percent. An * means the value was less than 1%.**

When the total site number and subplot numbers do not match between categories it is due to a portion the sites having been sampled in 1988 (sites 1-39) before those variables were being collected, or that particular data point was overlooked in the field. The latter explanation accounts for less than 1% of the occurrences.

DOMINANT SHRUB MEAN COVERAGE AND CONSTANCIES

Table SHRUB-1. Eastside lake RMZs, water type 1, dominant shrub #1 mean subplot coverage and constancy (total sites = 1, total subplots = 58). * value was less than 1.0
note: values are in percent

<u>Shrub Name</u>	<u>Coverage</u>	<u>Constancy</u>
snowberry	24	66
not present		16
bearberry	1	10
Douglas fir	*	2
water birch	*	2
russet buffaloberry	*	2
rose spp.	*	2
bristly Nootka rose	*	2

Table SHRUB-2. Eastside lake RMZs, water type 2, dominant shrub #1 mean subplot coverage and constancy (total sites = 1, total subplots = 37).

<u>Shrub Name</u>	<u>Coverage</u>	<u>Constancy</u>
not present		22
mallow ninebark	7	16
snowberry	4	14
ocean-spray	3	14
thimbleberry	1	11
hardhack	1	8
rose spp.	*	5
black hawthorne	*	3
baldhip rose	*	3
willow spp.	*	3
Douglas maple	1	3

Snowberry, bearberry, and mallow ninebark were the most frequently encountered dominant shrubs within eastside, water type 1 and 2, lake RMZs. It was not uncommon for shrubs to be lacking completely (i.e., not present).

Table SHRUB-3. Eastside lake RMZs, water type 1, dominant shrub #2 mean subplot coverage and constancy (total sites = 1, total subplots = 58). * value was less than 1.0
note: values are given in percent

<u>Shrub Name</u>	<u>Coverage</u>	<u>Constancy</u>
bearberry	1	12
russet buffaloberry	*	10
Douglas fir	*	5
snowberry	*	5
bristly Nootka rose	*	3
prickly currant	*	3
serviceberry	*	3
currant spp.	*	3
water birch	*	2
baldhip rose	*	2
huckleberry spp.	*	2
hardhack	*	2

Bearberry and russet buffaloberry were the most common sub-dominant shrubs within eastside, water type 1, lake RMZs.

Table SHRUB-4. Westside lakes, water type 1, dominant shrub #1 mean subplot cover age and constancy (total sites = 5, total subplots = 192). * value was less than 1.0 note: values are in percent

<u>Shrub Name</u>	<u>Coverage</u>	<u>Constancy</u>
salmonberry	13	28
salal	15	19
red huckleberry	1	9
not present		6
devil's club	2	6
Alaskan huckleberry	1	6
rusty menziesia	1	5
trailing blackberry	2	5
vine maple	3	5
Oregon grape	1	3
Indian plum	*	2
hardhack	1	2
ocean-spray	*	1
hazelnut	*	1
red-osier dogwood	*	1
baldhip rose	*	1
western hemlock	*	1
black cottonwood	*	1
stink currant	*	1

Table SHRUB-5. Westside lakes, water type 2, dominant shrub #1 mean subplot coverage and constancy (total sites = 3, total subplots = 129).

<u>Shrub Name</u>	<u>Coverage</u>	<u>Constancy</u>
salal	18	35
hardhack	6	12
salmonberry	8	12
trailing blackberry	2	12
Cascade Oregon grape	3	8
red huckleberry	1	6
Pacific ninebark	2	3
not present		2
evergreen huckleberry	*	2
rose spp.	*	2
snowberry	*	2
ocean-spray	*	1
red elderberry	*	1
Douglas fir	*	1
hazelnut	*	1

Table SHRUB-6. Westside lakes, water type 3, dominant shrub #1 mean subplot coverage and constancy (total sites = 1, total subplots = 72).

<u>Shrub Name</u>	<u>Coverage</u>	<u>Constancy</u>
salal	38	68
hardhack	25	32

Salmonberry, salal, and hardhack were the most commonly encountered dominant shrubs within westside, water type 1, 2, and 3 lake RMZs.

Table SHRUB-7. Westside lake RMZs, water type 1, dominant shrub #2 mean subplot coverage and constancy (total sites = 4, total subplots = 143). * value was less than 1.0
note: values are in percent

<u>Shrub Name</u>	<u>Coverage</u>	<u>Constancy</u>
not present		21
salmonberry	1	10
trailing blackberry	*	8
red huckleberry	*	8
rusty menziesia	*	7
Alaska huckleberry	*	7
Cascade Oregon grape	*	6
salal	1	4
Pacific ninebark	*	4
western hemlock	*	4
vine maple	*	4
hardhack	*	4
devil's club	*	3
serviceberry	*	2
oceanspray	*	2
Utah honeysuckle	*	1
red-osier dogwood	*	1
western red cedar	*	1
Indian plum	*	1
stink currant	*	1

Table SHRUB-8. Westside lake RMZs, water type 2, dominant shrub #2 mean subplot coverage and constancy (total sites = 2, total subplots = 67).

<u>Shrub Name</u>	<u>Coverage</u>	<u>Constancy</u>
not present		28
trailing blackberry	2	18
salal	1	13
red huckleberry	1	10
hardhack	2	9
Cascade Oregon grape	*	5
evergreen huckleberry	*	5
Pacific ninebark	2	3
baldhip rose	*	3
salmonberry	*	3
western hemlock	*	2
alder spp.	*	2

Table SHRUB-9. Westside Lake RMZs, water type 3, dominant shrub #2 mean subplot coverage and constancy (total sites = 1, total subplots = 73).

<u>Shrub Name</u>	<u>Coverage</u>	<u>Constancy</u>
not present		59
hardhack	2	18
salal	1	12
red huckleberry	*	3
trailing blackberry	*	3
western crabapple	1	3
unknown	*	1
western hemlock	*	1

Thirty-six percent of the time presence of a sub-dominant shrub in westside, water type 1, 2, and 3, lake RMZs was lacking. Salal, hardhack, salmonberry, and trailing blackberry were the most frequently encountered sub-dominant shrubs.

Table SHRUB-10. Eastside, boulder/bedrock, water type 3, dominant shrub #1 mean subplot coverage and constancy (total sites = 1, total subplots = 157). * value was less than 1.0 note: values are in percent

<u>Shrub Name</u>	<u>Coverage</u>	<u>Constancy</u>
big huckleberry	6	24
alder spp.	16	22
devil's club	5	13
rusty menziesia	5	12
willow spp.	7	8
stink currant	1	8
thimbleberry	2	5
salmonberry	1	3
mountain ash	*	3
prickly currant	*	1
pachistima	*	1
vine maple	*	1
not present		1

One eastside, boulder/bedrock, water type 1 RMZ was sampled in 1988. The most common dominant shrubs were big huckleberry, alder species and devil's club.

Table SHRUB-11. Westside, boulder/bedrock, water type 1, dominant shrub #1 mean subplot coverage and constancy (total sites = 7, total subplots = 522). * value was less than 1.0 note: values are in percent

Shrub Name	Coverage	Constancy
salmonberry	20	40
vine maple	16	25
not present		7
Alaska huckleberry	1	4
red huckleberry	*	4
salal	1	3
stink currant	*	3
trailing blackberry	*	3
red elderberry	*	2
devil's club	*	2
rusty menziesia	*	1
Pacific ninebark	*	1
Cascade Oregon grape	*	1
snowberry	*	1
Indian plum	*	1
red-osier dogwood	*	1
thimbleberry	*	1
hazelnut	*	1
western hemlock	*	1
western red cedar	*	1

Table SHRUB-12. Westside, boulder/bedrock, water type 2, dominant shrub #1 mean subplot coverage and constancy (total sites = 1, total subplots = 95).

<u>Shrub Name</u>	<u>Coverage</u>	<u>Constancy</u>
not present		23
red huckleberry	1	12
salal	4	12
Cascade Oregon grape	3	11
Alaska huckleberry	1	7
oceanspray	2	7
devil's club	1	6
vine maple	3	6
salmonberry	3	5
trailing blackberry	*	3
pachistima	*	2
big huckleberry	*	2
stink currant	*	1
baldhip rose	*	1
red elderberry	*	1

Table SHRUB-13. Westside, boulder/bedrock, water type 3, dominant shrub #1 mean subplot coverage and constancy (total sites = 2, total subplots = 115).

<u>Shrub Name</u>	<u>Coverage</u>	<u>Constancy</u>
salmonberry	25	44
not present		17
red huckleberry	1	7
salal	1	6
vine maple	2	5
trailing blackberry	*	5
stink currant	1	4
oceanspray	*	4
Cascade Oregon grape	*	3
western hemlock	*	2
devil's club	*	2
Indian plum	*	1
thimbleberry	*	1

Sixteen percent of the time presence of shrubs in westside, water type 1, 2, and 3, boulder/bedrock RMZs were lacking. When shrubs were encountered they were most frequently salmonberry, vine maple, red huckleberry and salal.

Table SHRUB-14. Westside, boulder/bedrock, water type 1, dominant shrub #2 mean subplot coverage and constancy (total sites = 7, total subplots = 334). * value was less than 1.0 note: values are in percent

<u>Shrub name</u>	<u>Coverage</u>	<u>Constancy</u>
not present		37
salmonberry	2	11
red huckleberry	1	8
stink currant	*	6
vine maple	*	6
salal	*	4
trailing blackberry	*	4
red elderberry	*	3
Alaska huckleberry	*	3
thimbleberry	*	3
devil's club	*	2
Indian plum	*	2
prickly currant	*	1
baldhip rose	*	1
western red cedar	*	1
snowberry	*	1
Pacific ninebark	*	1
western hemlock	*	1
Cascade Oregon grape	*	1
rusty menziesia	*	1

Table SHRUB-15. Westside, boulder/bedrock, water type 2, dominant shrub #2 mean subplot coverage and constancy (total sites = 1, total subplots = 51).

<u>Shrub name</u>	<u>Coverage</u>	<u>Constancy</u>
not present		49
salmonberry	*	8
pachistima	*	8
devil's club	2	8
Alaska huckleberry	*	6
red huckleberry	*	6
stink currant	*	6
Cascade Oregon grape	*	2
vine maple	1	2
salal	*	2
western hemlock	*	2
red elderberry	*	2

Table SHRUB-16. Westside, boulder/bedrock, water type 3, dominant shrub #2 mean subplot coverage and constancy (total sites = 2, total subplots = 83).

<u>Shrub name</u>	<u>Coverage</u>	<u>Constancy</u>
not present		53
stink currant	1	8
trailing blackberry	*	6
salmonberry	*	5
salal	*	5
western hemlock	*	4
red huckleberry	*	4
devil's club	1	4
vine maple	*	2
prickly currant	*	2
willow spp.	*	1
Cascade Oregon grape	*	1
twinfleur	*	1
douglas fir	*	1
big huckleberry	*	1
red elderberry	*	1

Sub-dominant shrubs were lacking in westside, boulder/bedrock, water type 1, 2, and 3 RMZs. When shrubs were found they most frequently were salmonberry, stink currant, pachistima, and devil's club.

Table SHRUB-17. Eastside, gravel/cobble, water type 1, dominant shrub #1 mean subplot coverage and constancy (total sites = 1, total subplots = 44). * value was less than 1.0 note: values are in percent

<u>Shrub Name</u>	<u>Coverage</u>	<u>Constancy</u>
snowberry	25	43
alder spp.	26	32
red-osier dogwood	4	5
mallow ninebark	*	2
mockorange	*	2
serviceberry	*	2
shiny leaf spirea	*	2
unknown	*	2
bittercherry	*	2
willow spp.	*	2
Douglas maple	13	2
not present		2

Table SHRUB-18. Eastside, gravel/cobble, water type 2, dominant shrub #1 mean coverage and constancy (total sites = 1, total subplots = 87).

<u>Shrub Name</u>	<u>Coverage</u>	<u>Constancy</u>
snowberry	13	28
willow spp.	4	18
red-osier dogwood	10	16
alder spp.	7	10
not present		8
baldhip rose	*	5
black hawthorne	*	3
rose spp.	*	2
mockorange	*	2
ocean-spray	*	2
Douglas maple	*	1
Cascade Oregon grape	*	1

Table SHRUB-19. Eastside, gravel/cobble, water type 3, dominant shrub #1 mean subplot coverage and constancy (total sites = 11, total subplots = 701).

<u>Shrub Name</u>	<u>Coverage</u>	<u>Constancy</u>
snowberry	8	18
alder spp.	9	17
red-osier dogwood	4	8
not present		8
vine maple	3	7
Douglas maple	3	6
thimbleberry	2	5
hazelnut	2	4
stink currant	*	3
prickly currant	*	3
pachistima	*	2
mockorange	1	2
devil's club	*	2
black hawthorne	*	2
baldhip rose	*	1
serviceberry	*	1
salmonberry	*	1
Cascade Oregon grape	*	1
Oregon grape	*	1
big huckleberry	*	1

Snowberry, alder species, willow species and red osier dogwood were the most frequently encountered dominant shrubs within eastside, gravel/cobble, water type 1, 2, and 3 RMZs.

Table SHRUB-20. Eastside, gravel/cobble, water type 1, dominant shrub #2 mean subplot coverage and constancy (total sites = 1, total subplots = 44). * value was less than 1.0 note: values are in percent

<u>Shrub Name</u>	<u>Coverage</u>	<u>Constancy</u>
snowberry	5	23
serviceberry	3	14
not present		9
poison-ivy	1	7
mockorange	1	7
alder spp.	3	7
bristly Nootka rose	1	5
unknown	*	5
ocean-spray	*	5
red-osier dogwood	1	5
willow spp.	*	2
Oregon grape	*	2
Douglas fir	*	2
bittercherry	1	2
thimbleberry	*	2
mallow ninebark	*	2

Table SHRUB-21. Eastside, gravel/cobble, water type 3, dominant shrub #2 mean subplot coverage and constancy (total sites = 7, total subplots = 425).

<u>Shrub Name</u>	<u>Coverage</u>	<u>Constancy</u>
snowberry	5	18
not present		17
thimbleberry	2	9
red-osier dogwood	1	7
Douglas maple	1	7
prickly currant	*	6
alder spp.	1	5
pachistima	1	5
mockorange	1	4
serviceberry	*	3
rose spp.	*	3
shiny leaf spirea	*	3
hazelnut	*	2
bristly Nootka rose	*	2
twinflor	*	1
mallow ninebark	*	1
blackcap	*	1
Oregon grape	*	1
rubus spp.	*	1
unknown	*	1

Snowberry, serviceberry and thimbleberry were the most frequently encountered sub-dominant shrub species within eastside, gravel/cobble, water type 1 and 3 RMZs. Water type 3 RMZs had a high percentage of subplots lacking in a sub-dominant shrub species.

Table SHRUB-22. Westside, gravel/cobble, water type 1, dominant shrub #1 mean subplot coverage and constancy (total sites = 10, total subplots = 892). * value was less than 1.0 note: values are in percent

<u>Shrub Name</u>	<u>Coverage</u>	<u>Constancy</u>
salmonberry	14	22
vine maple	15	20
red-osier dogwood	7	9
not present		9
Pacific ninebark	4	5
Alaska huckleberry	1	5
red huckleberry	*	3
red elderberry	*	3
Cascade Oregon grape	*	3
salal	1	2
Indian plum	1	2
trailing blackberry	*	2
rusty menziesia	*	2
alder spp.	*	1
mallow ninebark	*	1
devil's club	*	1
snowberry	*	1
big huckleberry	*	1
willow spp.	*	1
pachistima	*	1

Table SHRUB-23. Westside, gravel/cobble, water type 2, dominant shrub #1 mean subplot coverage and constancy (total sites = 8, total subplots = 704).

<u>Shrub Name</u>	<u>Coverage</u>	<u>Constancy</u>
salmonberry	31	48
salal	7	12
vine maple	7	11
red huckleberry	*	4
not present		4
trailing blackberry	1	4
Alaska huckleberry	1	3
devil's club	1	3
rusty menziesia	*	1
Pacific ninebark	*	1
cascara	*	1
hardhack	*	1
mallow ninebark	*	1
Indian plum	*	1
red elderberry	*	1
stink currant	*	1
black twinberry	*	1
Cascade Oregon grape	*	1
Utah honeysuckle	*	1
thimbleberry	*	1

Table SHRUB-24. Westside, gravel/cobble, water type 3, dominant shrub #1 mean subplot coverage and constancy (total sites = 39, total subplots = 3306).

<u>Shrub Name</u>	<u>Coverage</u>	<u>Constancy</u>
salmonberry	17	34
vine maple	12	18
salal	4	9
not present		9
trailing blackberry	1	4
devil's club	1	4
red elderberry	*	4
stink currant	1	4
red huckleberry	*	3
Cascade Oregon grape	*	2
red-osier dogwood	1	2
Alaska huckleberry	*	2
rusty menziesia	*	1
Indian plum	*	1
cascara	*	1
blackcap	*	1
Pacific ninebark	*	1
western hemlock	*	1
black twin-berry	*	1
big huckleberry	*	1

Salmonberry, salal and vine maple were the most common dominant shrub species within westside, gravel/cobble, water type 1, 2, and 3 RMZs.

Table SHRUB-25. Westside, gravel/cobble, water type 1, dominant shrub #2 mean subplot coverage and constancy (total sites = 10, total subplots = 832). * value was less than 1.0 note: values are in percent

<u>Shrub Name</u>	<u>Coverage</u>	<u>Constancy</u>
not present		22
salmonberry	4	14
vine maple	2	9
Indian plum	1	5
red elderberry	*	5
devil's club	1	5
red-osier dogwood	1	4
trailing blackberry	*	4
Cascade Oregon grape	*	3
snowberry	*	3
Pacific ninebark	*	3
rusty menziesia	*	3
red huckleberry	*	3
stink currant	*	2
Alaska huckleberry	*	2
big huckleberry	*	1
salal	*	1
twinflower	*	1
baldhip rose	*	1
western hemlock	*	

Table SHRUB-26. Westside, gravel/cobble, water type 2, dominant shrub #2 mean sub-plot coverage and constancy (total sites = 8, total subplots = 412).

<u>Shrub Name</u>	<u>Coverage</u>	<u>Constancy</u>
not present		29
salmonberry	3	17
Alaska huckleberry	1	9
red huckleberry	*	9
salal	*	7
vine maple	1	6
trailing blackberry	*	3
rusty menziesia	*	2
devil's club	*	2
red elderberry	*	2
western hemlock	*	2
stink currant	*	2
Pacific ninebark	*	2
blackcap	*	1
Cascade Oregon grape	*	1
hardhack	*	1
big huckleberry	*	1
Utah honeysuckle	*	1
red alder	*	1
alder spp.	*	1

Table SHRUB-27. Westside, gravel/cobble, water type 3, dominant shrub #2 mean subplot coverage and constancy (total sites = 39, total subplots = 2733).

<u>Shrub Name</u>	<u>Coverage</u>	<u>Constancy</u>
not present		35
salmonberry	2	13
stink currant	1	7
vine maple	1	6
red elderberry	*	5
devil's club	1	5
red huckleberry	*	5
trailing blackberry	*	5
salal	*	3
Alaska huckleberry	*	3
Cascade Oregon grape	*	2
western hemlock	*	2
Indian plum	*	1
rusty menziesia	*	1
red-osier dogwood	*	1
big huckleberry	*	1
casara	*	1
Pacific ninebark	*	1
blackcap	*	1
thimbleberry	*	1

Twenty-nine percent of the time westside, gravel/cobble, water type 1, 2, and 3 RMZs sampled lacked sub-dominant shrubs. When sub-dominant shrubs were present they were most frequently salmonberry, Alaskan huckleberry, vine maple and stink currant.

DOMINANT HERB MEAN COVERAGE AND CONSTANCIES

Table HERB-1. Eastside lakes, water type 1, dominant herb #1 mean subplot coverage and constancy (total sites = 1, total subplots = 58). * value was less than 1.0 note: values are given in percent

<u>Herb Name</u>	<u>Coverage</u>	<u>Constancy</u>
grass	31	69
pinegrass	3	5
horsetail	2	5
soft rush	*	4
rush	1	4
aster	*	2
lady-fern	*	2
Carex	*	2
Canada thistle	1	2
daisy	1	2
white flowered hawkweed	*	2
starry solomon	*	2
unknown	1	2

The most common dominant herbs within eastside, water type 1, lake RMZs were grass species, pine grass, and horsetails.

Table HERB-2. Eastside lakes, water type 1, dominant herb #2 mean subplot coverage and constancy (total sites = 1, total subplots = 58). * value was less than 1.0 note: values are given in percent

<u>Herb Name</u>	<u>Coverage</u>	<u>Constancy</u>
common yarrow	1	24
grass	2	14
unknown	1	10
not present		7
strawberry	*	7
thistle spp.	*	5
daisy	*	5
Carex	*	4
Canada thistle	*	4
soft rush	*	4
lupin	*	4
starry solomon	*	4
aster	*	2
fireweed	*	2
rush spp.	*	2
Mountain sweet root	*	2
buttercup	*	2
dock	1	2

The most commonly encountered sub-dominant herbs within eastside, water type 1, lake RMZs were common yarrow, grass species, and unknown species.

Table HERB-3. Westside lakes, water type 1, dominant herb #1 mean subplot coverage and constancy (total sites = 4, total subplots = 143). * value was less than 1.0 note: values are given in percent

<u>Herb Name</u>	<u>Coverage</u>	<u>Constancy</u>
swordfern	4	28
not present		23
lady-fern	1	10
wood-fern	*	9
deer-fern	*	8
bracken-fern	*	7
false lily of the valley	*	3
piggyback plant	2	3
goatsbeard	*	1
bunchberry dogwood	*	1
sweetscented bedstraw	*	1
carex spp.	*	1
rattlesnake plantain	*	1
grass	*	1
candy flower	*	1
licorice-fern	*	1
coolwort foamflower	*	1
western starflower	*	1
common cat-tail	*	1

Table HERB-4. Westside lakes, water type 2, dominant herb #1 mean subplot coverage and constancy (total sites = 2, total subplots = 67).

<u>Herb Name</u>	<u>Coverage</u>	<u>Constancy</u>
swordfern	13	46
not present		19
bracken-fern	1	11
tansy	*	6
deer-fern	*	5
vanilla leaf	1	3
wild ginger	*	2
lady-fern	*	2
carex spp.	*	2
fireweed	*	2
sweetscented bedstraw	*	2
grass	*	2
unknown	*	2

Table HERB-5. Westside lakes, water type 3, dominant herb #1 mean subplot coverage and constancy (total sites = 1, total subplots = 73).

<u>Herb Name</u>	<u>Coverage</u>	<u>Constancy</u>
not present		38
carex spp.	10	37
bracken-fern	1	19
grass	1	3
lady-fern	*	1
false lily of the valley	*	1

Swordfern and carex species were the most frequently encountered dominant herbs within westside, water type 1, 2, and 3, lake RMZs. The absence of herbs altogether was also common.

Table HERB-6. Westside lakes, water type 1, dominant herb #2, mean subplot coverage and constancy (total sites = 4, total subplots = 143). * value was less than 1.0
note: values are given in percent

<u>Herb Name</u>	<u>Coverage</u>	<u>Constancy</u>
not present		57
deer-fern	*	9
lady-fern	*	7
wood-fern	*	5
swordfern	*	5
bunchberry dogwood	*	2
false lily of the valley	*	2
goatsbeard	*	1
oak-fern	*	1
licorice-fern	*	1
bracken-fern	*	1
maidenhair-fern	*	1
fireweed	*	1
sweetscented bedstraw	*	1
grass	*	1
skunk cabbage	*	1
stag's horn moss	*	1
Cooley's hedgenettle	*	1
dandelion	*	1
western starflower	*	1

Table HERB-7. Westside lakes, water type 2, dominant herb #2, mean subplot coverage and constancy (total sites = 2, total subplots = 67).

<u>Herb Name</u>	<u>Coverage</u>	<u>Constancy</u>
not present		46
grass	*	11
bracken-fern	*	9
false lily of the valley	*	6
swordfern	*	6
wild ginger	*	5
lady-fern	*	5
tansy	*	5
vanilla leaf	*	2
deer-fern	*	2
fireweed	*	2
sweetscented bedstraw	*	2
white flowered hawkweed	*	2
Cooleye's hedgenettle	*	2

Table HERB-8. Westside lakes, water type 3, dominant herb #2, mean subplot coverage and constancy (total sites = 1, subplots = 73).

<u>Herb Name</u>	<u>Coverage</u>	<u>Constancy</u>
not present		71
false lily of the valley	*	18
western starflower	*	6
unknown	*	4
carex spp.	*	1

The majority of the time there was not a sub-dominant herb within westside, water type 1, 2, and 3, lake RMZs. When herbs were found they most frequently were grass species, false lily of the valley, and deer-fern.

Table HERB-9. Westside, boulder/bedrock, water type 1, dominant herb #1 mean sub-plot coverage and constancy (total sites = 7, total subplots = 334). * value was less than 1.0 note: values are given in percent

<u>Herb Name</u>	<u>Coverage</u>	<u>Constancy</u>
swordfern	17	44
Oregon oxalis	7	19
piggyback plant	1	4
grass	*	4
waterleaf	2	4
not present		3
lady-fern	1	3
deer-fern	*	3
coolwort foamflower	1	3
Scouler's corydalis	*	2
skunk cabbage	*	1
bunchberry dogwood	*	1
wood-fern	*	1
cow parsnip	*	1
wall lettuce	*	1
bracken-fern	*	1
Cooley's hedgenettle	*	1
goatsbeard	*	1
daisy	*	1
coltsfoot	*	1

Table HERB-10. Westside, boulder/bedrock, water type 2, dominant herb #1 mean sub-plot coverage and constancy (total sites = 2, total subplots = 51).

<u>Herb Name</u>	<u>Coverage</u>	<u>Constancy</u>
swordfern	34	75
not present		10
deer-fern	*	4
unknown	*	4
lady-fern	*	2
Scouler's corydalis	1	2
oak-fern	*	2
candy flower	*	2

Table HERB-11. Westside, boulder/bedrock, water type 3, dominant herb #1 mean sub-plot coverage and constancy (total sites = 2, total subplots = 83).

<u>Herb Name</u>	<u>Coverage</u>	<u>Constancy</u>
swordfern	20	54
piggyback plant	16	28
not present		4
deer-fern	*	4
bracken-fern	2	4
grass	*	1
candy flower	*	1
tansy	*	1
fringecup	*	1
trillium	*	1
unknown	*	1

The most commonly encountered dominant herbs within westside, boulder/bedrock, water type 1, 2, and 3, RMZs were swordfern, piggyback plant, and Oregon oxalis.

Table HERB-12. Westside, boulder/bedrock, water type 1, dominant herb #2 mean subplot coverage and constancy (total sites = 10, total subplots = 334). * value was less than 1.0 note: values are given in percent

<u>Herb Name</u>	<u>Coverage</u>	<u>Constancy</u>
not present		21
swordfern	1	12
Oregon oxalis	1	10
piggyback plant	1	9
grass	*	6
lady-fern	*	5
wood-fern	*	5
deer fern	*	4
waterleaf	*	3
coolwort foamflower	*	3
skunk cabbage	*	2
licorice-fern	*	2
sweetscented bedstraw	*	2
stinging nettle	*	2
horsetail	*	2
goatsbeard	*	1
false lilly of the valley	*	1
Scouler's corydalis	*	1
alumroot	*	1
unknown	*	1

Table HERB-13. Westside, boulder/bedrock, water type 2, dominant herb #2 mean subplot coverage and constancy (total sites = 1, total subplots = 51).

<u>Herb Name</u>	<u>Coverage</u>	<u>Constancy</u>
not present		26
deer-fern	1	18
trillium	*	12
wood-fern	*	10
lady-fern	*	8
Scouler's corydalis	*	8
swordfern	*	6
goatsbeard	*	4
unknown	*	4

Table HERB-14. Westside, boulder/bedrock, water type 3, dominant herb #2 mean subplot coverage and constancy (total sites = 2, total subplots = 83).

<u>Herb Name</u>	<u>Coverage</u>	<u>Constancy</u>
lady-fern	2	17
not present		16
swordfern	1	13
deer-fern	*	8
Scouler's corydalis	1	7
bracken-fern	*	7
dwarf nightshade	*	5
piggyback plant	1	5
sweetscented bedstraw	*	4
maidenhair-fern	*	2
grass	*	2
unknown	*	2
Columbia brome	*	1
wood-fern	*	1
horsetail	*	1
waterleaf	*	1
candyflower	*	1
licorice-fern	*	1
coolwort foamflower	*	1

Within westside, boulder/bedrock, water type 1, 2, and 3, RMZs it was not uncommon to find subdominant herbs lacking. When sub-dominant herbs were present they were most frequently swordfern, lady-fern, and deer-fern.

Table HERB-15. Eastside, gravel/cobble, water type 1, dominant herb #1 mean subplot coverage and constancy (total sites = 1, total subplots = 44). * value was less than 1.0
note: values are given in percent

<u>Herb Name</u>	<u>Coverage</u>	<u>Constancy</u>
grass	26	39
horsetail	5	25
not present		16
showy aster	*	2
strawberry spp.	*	2
northern bedstraw	*	2
cow parsnip	*	2
soft rush	*	2
lupin spp.	*	2
canarygrass	1	2
claspleaf twistedstalk	*	2

Table HERB-16. Eastside, gravel/cobble, water type 3, dominant herb #1 mean subplot coverage and constancy (total sites = 6, total subplots = 425).

<u>Herb Name</u>	<u>Coverage</u>	<u>Constancy</u>
grass	10	19
coolwort foamflower	3	9
wild sasparilla	2	8
meadowrue	*	5
canarygrass	3	5
beadlily	*	5
starry solomon-plume	*	5
stinging nettle	1	5
not present		5
sweetscented bedstraw	1	4
horsetail	1	3
claspleaf twistedstald	*	3
unknown	*	2
bunchberry dogwood	*	2
dwarf nightshade	*	2
heart-leaf arnica	*	1
bromus spp.	1	1
mountain sweet-root	*	1
lady-fern	*	1
thistle spp.	*	1

Within eastside, gravel/cobble, water type 1 and 3, RMZs the most commonly encountered dominant herbs were grass species, horsetail, and coolwort foamflower.

Table HERB-17. Eastside, gravel/cobble, water type 1, dominant herb #2 mean subplot coverage and constancy (total sites = 1, total subplots = 44). * value was less than 1.0
note: values are given in percent

<u>Herb Name</u>	<u>Coverage</u>	<u>Constancy</u>
horsetail	8	30
not present		21
grass	4	14
sweetscented bedstraw	*	7
heart-leaf arnica	*	7
unknown	*	7
fireweed	*	5
pinegrass	*	2
broadpetal strawberry	*	2
rush spp.	*	2
bracken fern	*	2
pioneer violet	*	2

Table HERB-18. Eastside, gravel/cobble, water type 3, dominant herb #2 mean subplot coverage and constancy (total sites = 11, total subplots = 426).

<u>Herb Name</u>	<u>Coverage</u>	<u>Constancy</u>
not present		12
grass	1	9
beadlily	*	8
coolwort foamflower	*	6
unknown	*	6
starry solomon-plume	*	5
meadowrue	*	5
stinging nettle	*	5
western yarrow	*	4
sweetscented bedstraw	*	4
wild sasparilla	*	4
claspleaf twistedstalk	*	3
dwarf nightshade	*	3
horsetail	*	2
lady-fern	*	2
bunchberry dogwood	*	2
meadow goldenrod	*	2
broadpetal stawberry	*	1
silky lupine	*	1
mountain sweetroot	*	1

Within eastside, gravel/cobble, water type 1 and 3, grass species, beadlily and sweetscented bedstraw were the most commonly encountered sub-dominant herbs. RMZs Within water type 3 RMZs sub-dominant forbs were lacking.

Table HERB-19. Westside, gravel/cobble, water type 1, dominant herb #1 mean sub-plot coverage and constancy (total sites = 10, total subplots = 828). * value was less than 1.0 note: values are given in percent

<u>Herb Name</u>	<u>Coverage</u>	<u>Constancy</u>
swordfern	12	27
piggyback plant	8	17
canarygrass	5	7
Oregon oxalis	3	7
bunchberry dogwood	*	5
lady-fern	1	5
not present		3
carex spp.	2	3
Scouler's corydalis	1	3
grass	*	2
bracken-fern	*	2
ground ivy	1	2
stinging nettle	1	2
inside-out-flower	*	2
beadlily	*	1
wood-fern	*	1
false lily of the valley	*	1
unknown	*	1
fireweed	*	1
vanilla leaf	*	1

Table HERB-20. Westside, gravel/cobble, water type 2, dominant herb #1 mean subplot coverage and constancy (total sites = 8, total subplots = 413).

<u>Herb Name</u>	<u>Coverage</u>	<u>Constancy</u>
swordfern	7	25
deer-fern	1	16
Oregon oxalis	5	11
piggyback plant	5	11
lady-fern	1	10
grass	2	4
false lily of the valley	*	2
not present		2
carex spp.	1	2
alumroot	*	2
water parsley	1	2
buttercup	*	2
small fruited bullrush	1	2
coolwort foamflower	*	2
canarygrass	*	2
skunkcabbage	*	1
Colleye's hedgenettle	*	1
stinging nettle	*	1
horsetail	*	1
ground ivy	*	1

Table HERB-21. Westside, gravel cobble, water type 3, dominant herb #1 mean subplot coverage and constancy (total sites = 40, total subplots = 2734).

<u>Herb Name</u>	<u>Coverage</u>	<u>Constancy</u>
swordfern	13	29
piggyback plant	8	14
Oregon oxalis	6	12
lady-fern	2	6
deer-fern	*	5
canarygrass	3	5
grass	1	3
small fruited bulrush	2	3
carex spp.	1	2
skunk cabbage	*	2
buttercup	1	2
stinging nettle	*	2
bleeding heart	*	2
not present		1
waterleaf	*	1
false lily of the valley	*	1
sweetscented bedstraw	*	1
Scouler's corydalis	*	1
water parsley	*	1
candy flower	*	1

Within westside, gravel/cobble, water type 1, 2, and 3, RMZs the most common dominant herbs were swordfern, piggyback plant, Oregon oxalis and deer-fern.

Table HERB-22. Westside, gravel/cobble, water type 1, dominant herb #2 mean subplot coverage and constancy (total sites = 10, total subplots = 828). *value was less than 1.0 note: values are given in percent

<u>Herb Name</u>	<u>Coverage</u>	<u>Constancy</u>
not present		12
swordfern	1	9
lady-fern	1	8
piggyback plant	1	7
Oregon oxalis	2	7
grass	*	6
wood-fern	*	5
false lily of the valley	*	4
stinging nettle	*	4
carex spp.	*	3
Scouler's corydalis	*	3
bunchberry dogwood	*	2
sweetscented bedstraw	*	2
beadlily	*	2
horsetail	*	2
bleeding heart	*	2
bracken-fern	*	2
candy flower	*	2
vanilla leaf	*	1

Table HERB-23. Westside, gravel/cobble, water type 2, dominant herb #2 mean sub-plot coverage and constancy (total sites = 8, total subplots = 412).

<u>Herb Name</u>	<u>Coverage</u>	<u>Constancy</u>
not present		16
Oregon oxalis	1	10
swordfern	*	8
grass	1	8
piggyback plant	1	8
lady-fern	*	7
deer-fern	*	7
false lily of the valley	*	5
water parsley	*	3
carex spp.	*	3
sweet-scented bedstraw	*	3
coolwort foamflower	*	3
skunk cabbage	*	2
wood-fern	*	2
alumroot	*	2
pioneer violet	*	2
licorice-fern	*	1
buttercup	*	1
Cooley's hedgenettle	*	1
unknown	*	1

Table HERB-24. Westside, gravel/cobble, water type 3, dominant herb #2 mean sub-plot coverage and constancy (total sites = 39, total subplots = 2732).

<u>Herb Name</u>	<u>Coverage</u>	<u>Constancy</u>
lady-fern	2	12
swordfern	1	10
not present		9
piggyback plant	1	8
Oregon oxalis	1	7
grass	1	5
deer-fern	*	5
false lily of the valley	*	4
sweetscented bedstraw	*	3
skunk cabbage	*	3
stinging nettle	*	3
bleeding heart	*	3
candy flower	*	2
water parsely	*	2
unknown	*	2
Cooley's hedgenettle	*	2
waterleaf	*	2
Scouler's corydalis	*	1
wood-fern	*	1
horsetail	*	1

Within westside, gravel/cobble, water type 1, 2, and 3, RMZs the most frequently encountered sub-dominant herbs were swordfern, lady-fern, and Oregon oxalis. On water type 1 RMZs sub-dominant herbs were most frequently lacking.

MEAN COVERAGE AND CONSTANCY VALUES FOR OVERSTORY CANOPY, TOTAL SHRUBS, FORBS, AND GRAMINOIDS.

The following tables display the total overstory canopy closure, total shrub coverage, total forb coverage, and total grass coverage within the subplots. Site and subplot numbers are provided. Total subplot numbers were used to determine the mean coverages.

For example: Table COVER-1 is read as... within eastside, lakeside, water type 1 RMZs the mean subplot canopy coverage was 55%, mean total shrub coverage and constancies were 38% and 83% respectively, mean total forb coverage and constancies were 27% and 93% respectively, and mean total grass coverage was 60% and 100% respectively. Where N.A. appears in the column indicates that there were no sites sampled within the defined category.

Lakeside RMZs

Table COVER-1. Eastside Lake RMZ Mean Coverage/Constancy for Subplot Canopy, Total Shrubs, Forbs, and Graminoids. Note: Coverage values given are in percent			
WATER TYPE	1	2	3
Canopy	55%	76%	N.A.
Shrubs	38/83	36/78	N.A.
Forbs	27/93	26/95	N.A.
Grass	60/100	16/46	N.A.
Number of sites	1	1	N.A.
Number of subplots	58	37	N.A.

Table COVER-2. Westside Lake RMZ Mean Coverage/Constancy for Subplot Canopy, Total Shrubs, Forbs, and Graminoids. Note: Coverage values given are in percent

WATER TYPE	1	2	3
Canopy	90%	80%	47%
Shrubs	59/94	61/98	69/93
Forbs	31/82	30/81	9/44
Grass	14/17	19/40	28/39
Number of sites	5	3	1
Number of sub-plots	191	129	75

Subplot overstory canopy closure for eastside lake RMZs appeared less than westside lake RMZs. Shrub canopy coverage and frequency appeared greater within westside lake RMZs. Forb canopy coverage appeared higher within westside lake RMZs, yet forb frequency was lower than those found in eastside sites. Grass canopy coverage was higher within type 1 lake RMZs on the eastside and similar between state sides on type 2 lakes. Grass frequency was higher in eastside lake RMZs.

Boulder/bedrock RMZs

Table COVER-3. Eastside, Boulder/Bedrock, RMZ Mean Coverage/Constancy for Subplot Canopy, Total Shrubs, Forbs and Graminoids. Note: Coverage values given are in percent

WATER TYPE	1	2	3
Canopy	N.A.	N.A.	61%
Shrubs	N.A.	N.A.	58/98
Forbs	N.A.	N.A.	32/88
Grass	N.A.	N.A.	6/17
Number of sites	N.A.	N.A.	1
Number of sub-plots	N.A.	N.A.	157

Table COVER-4. Westside, Boulder/Bedrock, RMZ Mean Coverage/Constancy for Subplot Canopy, Total Shrubs, Forbs and Graminoids. Note: Coverage values given are in percent

WATER TYPE	1	2	3
Canopy	87%	93%	88%
Shrubs	59/92	40/77	53/83
Forbs	51/97	33/90	47/97
Grass	8/37	7/19	5/30
Number of sites	10	2	4
Number of sub-plots	522	96	115

No type 3 streams were sampled on the east side of the state. Means for westside type 1 and 2 streams can be found in table COVER-3 and table COVER 4. Within type 3 streams the westside had greater overstory canopy closure and greater forb canopy closure. Shrub and grass canopy coverage was nearly equivalent between westside and eastside sites. Aside from grasses occurring twice as often in westside sites the frequencies of these variables were similar.

Gravel/cobble RMZs

Table COVER-5. Eastside, Gravel/Cobble, RMZ Mean Coverage/Constancy for Subplot Canopy, Total Shrubs, Forbs and Graminoids. Note: Coverage values given are in percent

WATER TYPE	1	2	3
Canopy	69%	72%	74%
Shrubs	81/98	50/92	59/92
Forbs	28/89	16/32	37/93
Grass	56/77	36/75	32/63
Number of sites	1	1	11
Number of sub-plots	44	87	701

Table COVER-6. Westside, Gravel/Cobble, RMZ Mean Coverage/Constancy for Subplot Canopy, Total Shrubs, Forbs and Graminoids. Note: Coverage values given are in percent

WATER TYPE	1	2	3
Canopy	87%	82%	82%
Shrubs	73/88	67/96	59/90
Forbs	52/93	44/97	59/97
Grass	25/48	23/49	27/53
Number of sites	11	12	50
Number of sub-plots	916	704	3,309

Overstory canopy closure and forb coverage was greater in westside RMZs. Grass coverage was greater within eastside RMZs than within westside RMZs. Shrub coverage on westside type 1 streams was lower than eastside yet higher than the eastside on type 2 waters. Shrub coverage was similar within RMZs on type 3 streams. Grass and shrubs were more frequently found within eastside RMZs. Forbs were more frequently found within westside RMZs.

MEAN COVERAGE AND CONSTANCY VALUES FOR WATER, ROCK, SOIL, ORGANIC GROUND COVER (OGC), DOWNED WOOD 1 (DW1), DOWNED WOOD 2 (DW2), & DOWNED WOOD 3 (DW3).

The following tables display the coverage and constancy values for total water, rock, soil, and organic ground cover. The number of subplots sampled is provided in parenthesis next to the water type.

Water coverage is based on open water. Rock coverage is based on exposed rock, and soil coverage is based on exposed soil. Organic ground cover includes litter, duff, mosses, lichens, and fungi. Organic ground cover does not include the downed wood coverage.

Downed wood classes are based on the amount of decay the log exhibits. Downed wood 1 logs are recently fallen trees with tight bark. Downed wood 2 logs are beginning to decay on the outside, but still have a solid center. Downed wood 3 logs are decayed throughout.

Lakes

	Eastside			Westside		
Water Type	1 (58)	2 (37)	3	1 (191)	2 (129)	3 (75)
Water	0/0	0/0	N.A.	13/4	6/3	0/0
Rock	18/31	3/8	N.A.	4/9	3/2	3/1
Soil	8/33	3/22	N.A.	8/6	13/5	15/1
OGC	87/98	93/100	N.A.	92/99	96/99	97/89
	Eastside			Westside		
Water Type	1 (58)	2 (37)	3	1 (191)	2 (129)	3 (75)
DW1	3/2	7/38	N.A.	25/13	8/14	3/1
DW2	11/26	7/30	N.A.	9/22	7/23	5/7
DW3	8/38	13/27	N.A.	17/39	9/25	15/40

Boulder/bedrock

		Eastside			Westside		
Water Type	1	2	3 (157)	1 (522)	2 (95)	3 (116)	
Water	N.A.	N.A.	4/6	9/3	3/1	9/5	
Rock	N.A.	N.A.	15/33	18/31	24/27	6/32	
Soil	N.A.	N.A.	20/24	8/22	11/19	16/18	
OGC	N.A.	N.A.	81/98	86/99	87/99	92/99	
		Eastside			Westside		
Water Type	1	2	3 (157)	1 (522)	2 (95)	3 (116)	
DW1	N.A.	N.A.	10/31	8/13	9/2	10/22	
DW2	N.A.	N.A.	10/31	19/26	11/20	7/25	
DW3	N.A.	N.A.	13/20	17/31	19/40	7/28	

Gravel/cobble

		Eastside			Westside		
Water Type	1 (44)	2 (87)	3 (701)	1 (914)	2 (704)	3 (3306)	
Water	3/2	3/1	7/8	15/4	7/8	15/8	
Rock	23/30	26/62	5/14	18/9	10/4	9/7	
Soil	25/18	19/61	11/21	10/9	10/14	9/14	
OGC	82/96	68/99	86/99	93/96	93/99	93/99	
		Eastside			Westside		
Water Type	1 (44)	2 (87)	3 (701)	1 (914)	2 (704)	3 (3306)	
DW1	3/2	7/15	11/19	12/7	11/17	10/14	
DW2	12/9	4/12	11/26	11/16	11/14	12/20	
DW3	14/14	10/12	13/37	18/24	17/39	15/27	

LIVE TREE DENSITY

Tree diameter was measured in the following four inch size class intervals:

Size Class	Diameter in inches
1	0.0 - 3.9
2	4.0 - 7.9
3	8.0 - 11.9
4	12.0 - 15.9
5	16.0 - 19.9
6	20.0 - 23.9
7	24 +

Data were analyzed to determine the number of trees per acre and per 1000 feet within each size class. Size class analysis occurred on sizes 1-7, 2-7, 3-7, 4-7. When the last size class shown is 3-7, there were no trees larger than 11.9 inches in diameter within the defined category.

For example: Table TREE-1 is read as... there was a mean of seven conifers greater than 12.0 inches in diameter per 1000 feet within eastside lakes, water type 1. In this example a mean of seven trees per 1000 feet equates to a mean of eight trees (greater than 12.0 inches in diameter) per acre.

Trees analyzed as live fit one of the following criteria: live tree - undamaged, live tree - 1/3 to 1/2 of the top broken, live tree - dead top. Minimum height was 4.5 feet. All trees were grouped together by size class and category.

Trees were defined as either hardwood or conifer. The number of sites sampled and the total number of strips within these sites have been provided in the tables.

Strip count is not the total number of strips within the sampled RMZs, but instead is the total number of strips, in that category of RMZs, containing trees of the defined size class range. The total number of strips sampled within each RMZ category is not shown.

Trees/1000 feet and trees/per acre were calculated by dividing by the total number of trees (within the size class range) by the strip count.

The number of strips and sites decreased when trees no longer met the minimum size requirements. For example in Table TREE-9 (water type 3) the number of sites with trees in size classes 1-7 equals 9. The number of sites with trees in size classes 4-7 equals 7. This means there were two eastside gravel/cobble RMZs without trees larger than 12.0 inches in diameter. Strip count decreased from 135 to 87. Again, this means that 135 strips had at least one conifer within them, but only 87 strips had at least one conifer over 12.0 inches in diameter.

Lakeside Mean Tree Densities

WATER TYPE	SIZE CLASS	TREES/ ACRE	TREES/ 1000 FT	NUMBER OF SITES	NUMBER OF STRIPS
1	1-7	40	43	1	12
	2-7	29	31	1	12
	3-7	15	16	1	12
	4-7	8	7	1	10
2	1-7	51	42	1	10
	2-7	43	36	1	10
	3-7	24	20	1	10
	4-7	12	10	1	8

WATER TYPE	SIZE CLASS	TREES/ ACRE	TREES/ 1000 FT	NUMBER OF SITES	NUMBER OF STRIPS
1	1-7	7	8	1	4
	2-7	1	1	1	1
2	1-7	9	7	1	6
	2-7	8	6	1	5
	3-7	3	2	1	2

Table TREE-3. Westside Lake RMZ Mean Tree Density - Conifers

WATER TYPE	SIZE CLASS	TREES/ ACRE	TREES/ 1000 FT	NUMBER OF SITES	NUMBER OF STRIPS
1	1-7	42	41	5	26
	2-7	21	23	5	26
	3-7	11	13	5	25
	4-7	7	7	5	22
2	1-7	13	41	3	7
	2-7	9	15	3	6
	3-7	5	9	3	6
	4-7	3	6	3	6
3	1-7	25	30	1	9
	2-7	18	22	1	9
	3-7	8	10	1	8
	4-7	2	3	1	5

Table TREE-4. Westside Lake RMZ Mean Tree Density - Hardwoods

WATER TYPE	SIZE CLASS	TREES/ ACRE	TREES/ 1000 FT	NUMBER OF SITES	NUMBER OF STRIPS
1	1-7	37	88	5	18
	2-7	25	61	5	18
	3-7	17	44	5	16
	4-7	11	30	5	15
2	1-7	50	67	3	20
	2-7	42	54	3	20
	3-7	23	30	3	19
	4-7	7	10	3	15
3	1-7	13	15	1	10
	2-7	12	14	1	9
	3-7	5	6	1	6
	4-7	1	2	1	4

Statewide, type 1 lake RMZs contained similar amounts of conifers per 1000 feet and per acre. Conifer size in type 1 RMZs was similar on both sides of the state. Hardwood composition within water type 1 RMZs was considerably higher on the westside of the state. Conifers composition, within water type 2 RMZs, was higher in eastside sites. Tree size was larger in eastside RMZs. Hardwoods were more prevalent and larger in westside, type 2 RMZs.

Boulder/bedrock Mean Tree Densities

Table TREE-5. Eastside Boulder/Bedrock RMZ Mean Tree Density - Conifers

WATER TYPE	SIZE CLASS	TREES/ ACRE	TREES/ 1000 FT	NUMBER OF SITES	NUMBER OF STRIPS
3	1-7	27	18	1	42
	2-7	24	17	1	41
	3-7	12	8	1	33
	4-7	6	4	1	26

Table TREE-6. Eastside Boulder/Bedrock RMZ Mean Tree Density - Hardwoods

WATER TYPE	SIZE CLASS	TREES/ ACRE	TREES/ 1000 FT	NUMBER OF SITES	NUMBER OF STRIPS
3	1-7	2	1	1	8
	2-7	2	1	1	8
	3-7	1	1	1	7
	4-7	1	1	1	4

Table TREE-7. Westside Boulder/Bedrock RMZ Mean Tree Density - Conifers

WATER TYPE	SIZE CLASS	TREES/ACRE	TREES/1000 FT	NUMBER OF SITES	NUMBER OF STRIPS
1	1-7	30	54	10	54
	2-7	14	25	10	53
	3-7	7	13	10	47
	4-7	5	8	10	43
2	1-7	56	116	2	14
	2-7	42	85	2	13
	3-7	23	47	2	13
	4-7	12	8	2	11
3	1-7	60	41	4	24
	2-7	25	19	4	23
	3-7	11	10	4	19
	4-7	4	4	3	12

Table TREE-8. Westside Boulder/Bedrock RMZ Mean Tree Density - Hardwoods

WATER TYPE	SIZE CLASS	TREES/ACRE	TREES/1000 FT	NUMBER OF SITES	NUMBER OF STRIPS
1	1-7	42	68	10	78
	2-7	24	38	10	78
	3-7	14	21	10	70
	4-7	9	14	10	65
2	1-7	19	31	2	14
	2-7	14	23	2	14
	3-7	9	17	2	13
	4-7	5	10	2	12
3	1-7	54	42	4	26
	2-7	28	25	4	26
	3-7	16	16	4	21
	4-7	10	12	4	19

Westside, water type 3, boulder/bedrock RMZs had higher densities of hardwoods per acre and per 1000 feet. The composition of conifers between the two sides of the state were relatively equal. Within westside water type 1 RMZs hardwoods dominated over conifers. On water type 2 RMZs conifers dominated the hardwoods.

Gravel/cobble Mean Tree Densities

WATER TYPE	SIZE CLASS	TREES/ACRE	TREES/1000 FT	NUMBER OF SITES	NUMBER OF STRIPS
1	1-7	12	10	1	8
	2-7	10	8	1	8
	3-7	6	5	1	8
	4-7	4	4	1	8
2	1-7	9	5	1	9
	2-7	8	4	1	9
	3-7	4	3	1	8
	4-7	3	2	1	8
3	1-7	51	26	9	135
	2-7	20	14	9	129
	3-7	10	7	9	110
	4-7	7	5	7	87

WATER TYPE	SIZE CLASS	TREES/ACRE	TREES/1000 FT	NUMBER OF SITES	NUMBER OF STRIPS
1	1-7	10	8	1	7
	2-7	5	4	1	6
	3-7	3	3	1	4
	4-7	2	2	1	4
2	1-7	43	25	1	18
	2-7	30	17	1	18
	3-7	5	3	1	12
	4-7	3	2	1	7
3	1-7	29	25	11	115
	2-7	11	10	11	89
	3-7	4	4	11	71
	4-7	3	2	10	54

Table TREE-11. Westside Gravel/Cobble RMZ Mean Tree Density - Conifers

WATER TYPE	SIZE CLASS	TREES/ACRE	TREES/1000 FT	NUMBER OF SITES	NUMBER OF STRIPS
1	1-7	36	57	11	84
	2-7	12	19	11	76
	3-7	7	11	10	66
	4-7	4	7	10	60
2	1-7	35	40	9	77
	2-7	13	17	9	74
	3-7	9	11	9	66
	4-7	6	8	9	59
3	1-7	22	33	49	357
	2-7	11	16	47	332
	3-7	6	9	47	291
	4-7	4	6	46	243

Table TREE-12. Westside Gravel/Cobble RMZ Mean Tree Density - Hardwoods

WATER TYPE	SIZE CLASS	TREES/ACRE	TREES/1000 FT	NUMBER OF SITES	NUMBER OF STRIPS
1	1-7	35	81	11	115
	2-7	21	60	11	111
	3-7	16	42	11	106
	4-7	9	25	11	95
2	1-7	33	33	12	131
	2-7	27	27	11	120
	3-7	19	19	11	113
	4-7	12	13	11	107
3	1-7	31	60	50	495
	2-7	20	26	50	470
	3-7	14	17	50	431
	4-7	9	11	48	387

Westside, water type 1, gravel/cobble RMZs contained more conifers and hardwoods per acre than their eastside counterparts. Westside water type 2 RMZs contained more conifers per 1000 feet, and per acre than did their eastside counterparts, but fewer hardwoods. Westside water type 3 RMZs contained fewer conifers and more hardwoods per acre than similar eastside sites.

SNAG DENSITY

Snags were defined in the following manner: recent dead (needles or leaves dead, yet still on the tree), dead tree - tight bark, or dead tree - loose bark. Minimum height was 4.5 feet. There was no minimum size requirement for snags. All snags were grouped together by size class and category.

Lakeside Mean Snag Densities

WATER TYPE	SIZE CLASS	SNAGS/ ACRE	SNAGS/ 1000 FT	NUMBER OF SITES	NUMBER OF STRIPS
2	1-7	6	5	1	5
	2-7	5	4	1	5
	3-7	2	1	1	3
	4-7	1	1	1	1

WATER TYPE	SIZE CLASS	SNAGS/ ACRE	SNAGS/ 1000 FT	NUMBER OF SITES	NUMBER OF STRIPS
1	1-7	7	8	1	11
	2-7	4	5	1	9
	3-7	2	2	1	5
	4-7	2	2	1	4
2	1-7	3	3	1	4
	2-7	3	3	1	4
	3-7	1	1	1	2

Table SNAG-3. Westside Lake RMZ Mean Snag Density - Conifers

WATER TYPE	SIZE CLASS	SNAGS/ ACRE	SNAGS/ 1000 FT	NUMBER OF SITES	NUMBER OF STRIPS
1	1-7	5	5	4	15
	2-7	3	3	4	15
	3-7	3	3	3	13
	4-7	3	4	2	10
2	1-7	3	4	2	5
	2-7	1	2	2	3
	3-7	1	2	1	2
	4-7	1	1	1	1
3	1-7	18	21	1	9
	2-7	9	11	1	9
	3-7	2	3	1	6
	4-7	1	1	1	4

Table SNAG-4. Westside Lake RMZ Mean Snag Density - Hardwoods

WATER TYPE	SIZE CLASS	SNAGS/ ACRE	SNAGS/ 1000 FT	NUMBER OF SITES	NUMBER OF STRIPS
1	1-7	5	16	4	12
	2-7	5	15	3	9
	3-7	4	12	2	6
	4-7	2	5	2	4
2	1-7	11	14	3	17
	2-7	10	13	3	16
	3-7	5	6	3	13
	4-7	1	5	2	4
3	1-7	11	13	1	8
	2-7	7	8	1	7
	3-7	3	4	1	7
	4-7	2	5	2	4

Westside, water type 1, lake RMZs contained more hardwood snags per acre than conifers. Eastside, water type 2, RMZs contained more conifer, and similar hardwood snags per acre, than their westside counterparts. Westside, water type 3 RMZs contained more conifer snags per acre than hardwoods.

Boulder/bedrock Snag Densities

Table SNAG-5. Eastside, Boulder/Bedrock Mean Snag Density - Conifers

WATER TYPE	SIZE CLASS	SNAGS/ ACRE	SNAGS/ 1000 FT	NUMBER OF SITES	NUMBER OF STRIPS
3	1-7	4	3	1	17
	2-7	4	3	1	17
	3-7	2	2	1	15
	4-7	1	1	1	10

Table SNAG-6. Eastside, Boulder/Bedrock Mean Snag Density - Hardwoods

WATER TYPE	SIZE CLASS	SNAGS/ ACRE	SNAGS/ 1000 FT	NUMBER OF SITES	NUMBER OF STRIPS
3	1-7	1	1	1	2
	2-7	1	1	1	2
	3-7	1	1	1	2
	4-7	1	1	1	2

Table SNAG-7. Westside, Boulder/Bedrock Mean Snag Density - Conifers

WATER TYPE	SIZE CLASS	SNAGS/ ACRE	SNAGS/ 1000 FT	NUMBER OF SITES	NUMBER OF STRIPS
1	1-7	4	7	9	19
	2-7	2	3	9	18
	3-7	1	2	7	12
	4-7	1	2	6	11
2	1-7	12	26	2	9
	2-7	7	14	2	9
	3-7	2	3	2	6
	4-7	1	1	2	4
3	1-7	11	8	3	13
	2-7	4	3	3	9
	3-7	2	1	3	5
	4-7	1	1	2	3

Table SNAG-8. Westside, Boulder/Bedrock Mean Snag Density - Hardwoods

WATER TYPE	SIZE CLASS	SNAGS/ ACRE	SNAGS/ 1000 FT	NUMBER OF SITES	NUMBER OF STRIPS
1	1-7	4	7	10	35
	2-7	2	3	10	27
	3-7	1	2	10	17
	4-7	1	1	6	8
2	1-7	2	3	2	7
	2-7	2	3	2	6
	3-7	2	3	2	6
	4-7	2	2	2	6
3	1-7	5	4	4	13
	2-7	3	2	4	9
	3-7	2	2	4	5
	4-7	1	1	3	4

Westside, boulder/bedrock, water type 1, RMZs contained an equal ratio (per acre) of conifer to hardwood snags. Westside, type 2, RMZs contained more conifer snags per acre than hardwoods. Westside, type 3, RMZs contained more conifer and hardwood snags per acre than did their counterpart eastside sites. Westside and eastside, type 3, RMZs contained more conifer snags than hardwoods per acre.

Gravel/cobble Mean Snag Densities

Table SNAG-9. Eastside, Gravel/Cobble Mean Snag Density - Conifers

WATER TYPE	SIZE CLASS	SNAGS/ ACRE	SNAGS/ 1000 FT	NUMBER OF SITES	NUMBER OF STRIPS
1	1-7	1	1	1	1
2	1-7	3	4	1	6
	2-7	3	2	1	6
	3-7	1	2	1	5
	4-7	1	1	1	2
3	1-7	3	2	9	46
	2-7	1	1	8	32
	3-7	1	1	6	17
	4-7	1	1	6	14

Table SNAG-10. Eastside, Gravel/Cobble Mean Snag Density - Hardwoods

WATER TYPE	SIZE CLASS	SNAGS/ ACRE	SNAGS/ 1000 FT	NUMBER OF SITES	NUMBER OF STRIPS
1	1-7	1	1	1	2
	2-7	1	1	1	1
	3-7	1	1	1	1
	4-7	1	1	1	1
2	1-7	3	2	1	5
	2-7	2	1	1	5
	3-7	1	1	1	3
	4-7	1	1	1	1
3	1-7	7	6	8	50
	2-7	4	4	8	42
	3-7	2	2	5	23
	4-7	1	1	5	13

Table SNAG-11. Westside, Gravel/Cobble Mean Snag Density - Conifers

WATER TYPE	SIZE CLASS	SNAGS/ ACRE	SNAGS/ 1000 FT	NUMBER OF SITES	NUMBER OF STRIPS
1	1-7	3	5	9	33
	2-7	1	2	7	23
	3-7	1	1	4	11
	4-7	1	2	3	9
2	1-7	3	4	9	42
	2-7	2	2	9	34
	3-7	1	2	8	25
	4-7	1	1	6	18
3	1-7	2	5	43	153
	2-7	2	3	42	132
	3-7	1	2	35	80
	4-7	1	2	26	57

WATER TYPE	SIZE CLASS	SNAGS/ ACRE	SNAGS/ 1000 FT	NUMBER OF SITES	NUMBER OF STRIPS
1	1-7	4	18	10	60
	2-7	3	10	10	54
	3-7	1	2	6	28
	4-7	1	1	5	13
2	1-7	5	5	9	56
	2-7	4	4	9	49
	3-7	2	2	8	28
	4-7	1	1	8	16
3	1-7	5	7	43	250
	2-7	3	4	40	212
	3-7	2	2	35	122
	4-7	1	1	30	67

Westside, gravel/cobble, water type 1, RMZs contained more conifer, and fewer hardwood snags per acre, than eastside sites within the same category. Westside, water type 2, RMZs contained similar numbers of conifer snags, and more hardwood snags per acre, than do eastside, water type 2 RMZs. Within water type 3 RMZs there were similar numbers of conifer snags per acre between state sides, and more hardwood snags per acre in eastside RMZs. Hardwood snags dominated within all water types on both sides of the state.

UMAs

One hundred and twenty-six acres of UMAs located on 30 sites were sampled in 1988/89. UMAs were stratified by their dominant vegetative characteristics. The structure of the UMAs sampled in 1988/89 was a diverse array of forest types ranging from wetlands to old-growth forests.

UMAs are categorized by their physical characteristics. Three categories were developed: forested wetland, upland forest, and bogs. No bog UMAs were sampled on the eastside of the state.

VEGETATION AND OTHER STIRP VARIABLES

Data were collected on the two dominant shrubs and herbs, total shrubs, herbs and graminoids (grass), downed wood 1 to 3 (decay class 1 = recent fallen, decay class 3 = rotten), water, rock, and soil. Mean coverage and constancy were calculated on these variables

Canopy is defined as the percent of closed canopy above the sample plot. Coverage is defined as the percentage of ground, when viewed from above the subplot, the variable covers within the sample plot. Sample plots are 5x10 feet. Constancy is defined as the degree of presence a variable has within sample plots. Subplot coverage and constancy values are given in percent.

UMA shrubs and herbs are listed in order by their constancy values. Shrub tables 28 through 37 and herb tables 25 through 34 list the 20 most frequently encountered shrubs or forbs. When fewer than 20 shrubs or forbs are listed, this implies that fewer than 20 were encountered within that specific category.

When the total site number and subplot numbers do not match between categories it is because a portion of the sites, the first 39, were sampled in 1988 before those variables were being collected or that particular data point was overlooked in the field. The latter explanation accounts for less than 1% of the occurrences.

Values are given in percent. An * means the value was less than 1%.

DOMINANT SHRUB MEAN COVERAGE AND CONSTANCIES

Table SHRUB-28. Eastside UMAs, forested wetlands, dominant shrub #1 mean subplot coverage and constancy (total sites = 1, total subplots = 174).

<u>Shrub Name</u>	<u>Coverage</u>	<u>Constancy</u>
hardhack	9	21
snowberry	5	20
alder spp.	5	13
quaking aspen	*	10
not present		8
unknown	4	8
red-osier dogwood	1	6
prickly currant	*	4
devil's club	1	2
black hawthorne	1	2
western red cedar	*	2
thimbleberry	*	2
stink currant	*	1
baldhip rose	*	1
Oregon grape	*	1
serviceberry	*	1
Douglas maple	*	1

Within Eastside, forested wetland , UMAs the most commonly encountered dominant shrubs were hardhack, snowberry and alder species.

Table SHRUB-29. Eastside UMAs, upland forest, dominant shrub #1 mean subplot coverage and constancy (total sites = 2, total subplots = 197).

<u>Shrub Name</u>	<u>Covrage</u>	<u>Constancy</u>
mallow ninebark	19	27
not present		15
subalpine fir	*	10
low huckleberry	2	10
pachistima	1	8
twinflower	1	6
big huckleberry	1	6
baldhip rose	*	6
Utah honeysuckle	*	4
Douglas fir	*	4
common prince's pine	*	3
Oregon grape	*	3
unknown	*	1
grand fir	*	1

Within eastside, upland forest, UMAs the most common dominant shrubs (when present) were mallow ninebark and subalpine fir. The absence of a dominant shrub was recorded 15% of the time.

Table SHRUB-30. Eastside UMA, forested wetland, dominant shrub #2 mean subplot coverage and constancy (total sites = 1, total subplots = 174).

<u>Shrub Name</u>	<u>Coverage</u>	<u>Constancy</u>
not present		29
hardhack	2	11
quaking aspen	*	11
alder spp.	1	9
unknown	1	8
prickly currant	*	7
snowberry	1	6
red-osier dogwood	1	6
serviceberry	*	3
baldhip rose	*	3
thimbleberry	*	2
salmonberry	*	1
western red cedar	*	1
rubus spp.	*	1
Douglas maple	*	1
Oregon grape	*	1
rose spp.	*	1
devil's club	*	1
grand fir	*	1

Within eastside, forested wetland, UMAs a sub-dominant shrub was most frequently lacking. When sub-dominant shrubs were present they were most frequently hardhack and quaking aspen.

Table SHRUB-31. Eastside UMA, upland forest, dominant shrub #2 mean subplot coverage and constancy (total sites = 2, total subplots = 197).

<u>Shrub Name</u>	<u>Coverage</u>	<u>Constancy</u>
not present		28
snowberry	1	10
common prince's pine	*	9
pachistima	1	8
shiny leaf spirea	2	7
subalpine fir	*	7
big huckleberry	*	6
baldhip rose	*	6
low huckleberry	*	5
Douglas fir	*	5
Utah honeysuckle	*	5
Oregon grape	*	3
twinflor	*	3
Douglas fir	*	1
serviceberry	*	1
mallow ninebark	*	1

Within eastside, upland forest, UMAs sub-dominant shrub most frequently were lacking. When sub-dominant shrubs were encountered they were most frequently snowberry and common prince's pine.

Table SHRUB-32. Westside UMA, bogs, dominant shrub #1 mean subplot coverage and constancy (total sites = 2, total subplots = 273).

<u>Shrub Name</u>	<u>Coverage</u>	<u>Constancy</u>
salal	18	28
smooth Labrador-tea	16	22
western crabapple	13	17
hardback	12	14
vine maple	7	8
swamp laurel	4	7
casara	1	2
western hemlock	*	1
red huckleberry	*	1
not present		1

Within westside, bog, UMAs the most common dominant shrubs were salal, smooth Labrador-tea, and western crabapple.

Table SHRUB-33. Westside UMA, forested wetlands, dominant shrub #1 mean subplot coverage and constancy (total sites = 6, total subplots = 581).

<u>Shrub Name</u>	<u>Coverage</u>	<u>Constancy</u>
salmonberry	11	29
not present		21
vine maple	6	9
western crabapple	7	8
salal	3	6
red huckleberry	1	6
trailing blackberry	1	5
rusty menziesia	*	4
Alaska huckleberry	1	4
blackcap	*	1
big huckleberry	*	1
twinflower	*	1
red alder	*	1
western hemlock	*	1
red elderberry	*	1
devil's club	*	1
black twin-berry	*	1
cascara	*	1
Sitka spruce	*	1
sticky currant	*	1

Within westside, forested wetland, UMAs the most frequently encountered dominant shrubs were salmonberry and vine maple. Shrubs were lacking 21% of the time.

Table SHRUB-34. Westside UMA, upland forest, dominant shrub #1 mean subplot coverage and constancy (total sites = 11, total subplots = 1462).

<u>Shrub Name</u>	<u>Coverage</u>	<u>Constancy</u>
vine maple	13	20
salmonberry	5	12
trailing blackberry	3	8
not present		7
salal	2	7
Cascade Oregon grape	2	6
red huckleberry	1	6
hazelnut	2	5
big huckleberry	1	4
rusty menziesia	1	4
devil's club	1	2
red elderberry	*	2
baldhip rose	*	2
red-osier dogwood	1	1
ocean-spray	*	1
dwarf bramble	*	1
western hemlock	*	1
serviceberry	*	1
western red cedar	*	1
Himalayan blackberry	*	1

Within westside, upland forest, UMAs the most frequently encountered dominant shrubs were vine maple, salmonberry, and trailing blackberry.

Table SHRUB-35. Westside UMA, bog, dominant shrub #2 mean subplot coverage and constancy (total sites = 2, total subplot = 275).

<u>Shrub Name</u>	<u>Coverage</u>	<u>Constancy</u>
salal	11	25
western crabapple	6	15
hardhack	4	13
not present		12
swamp laurel	2	10
smooth Labrador-tea	2	10
vine maple	1	6
rusty menziesia	*	4
red huckleberry	*	2
salmonberry	*	2
cascara	*	1
western hemlock	*	1
Alaska huckleberry	*	1
Sitka spruce	*	1

Within westside, bog, UMAs the most commonly encountered sub-dominant shrubs were salal, western crabapple, and hardhack.

Table SHRUB-36. Westside UMA, forested wetlands, dominant shrub #2 mean subplot coverage and constancy (total sites = 6, total subplot = 551).

<u>Shrub Name</u>	<u>Coverage</u>	<u>Constancy</u>
not present		48
salmonberry	2	10
salal	2	9
red huckleberry	*	5
Alaska huckleberry	1	5
red huckleberry	*	5
rusty menziesia	*	4
trailing blackberry	*	3
western hemlock	*	3
vine maple	*	2
devil's club	*	1
stink currant	*	1
western crabapple	*	1
blackcap	*	1
Sitka spruce	*	1
big huckleberry	*	1
cascara	*	1
Pacific ninebark	*	1
hardhack	*	1
twinflower	*	1

Within westside, forested wetland, UMAs sub-dominant shrubs were predominantly lacking. When sub-dominant shrubs were present they most frequently were salmonberry and salal.

Table SHRUB-37. Westside UMA, upland forests, dominant shrub #2 mean subplot coverage and constancy (total sites = 11, total subplot = 886).

<u>Shrub Name</u>	<u>Coverage</u>	<u>Constancy</u>
not present		26
salmonberry	1	12
red huckleberry	*	7
Cascade Oregon grape	1	7
trailing blackberry	*	5
vine maple	1	5
rusty menziesia	*	5
salal	*	4
big huckleberry	*	4
red elderberry	*	3
western hemlock	*	2
common prince's pine	*	2
pachistima	*	2
Himalayan blackberry	*	2
baldhip rose	*	2
hazelnut	*	1
devil's club	*	1
dwarf bramble	*	1
unknown	*	1
thimbleberry	*	1

Within westside, upland forests, UMAs sub-dominant shrubs were generally lacking. When sub-dominant shrubs were found they most commonly were salmonberry and red huckleberry.

Dominant herb Mean Coverage and Constancies

Table HERB-25. Eastside UMAs, forested wetlands, dominant herb #1 mean subplot coverage and constancy (total sites = 1, total subplots = 197).

<u>Herb Name</u>	<u>Coverage</u>	<u>Constancy</u>
lady-fern	22	47
grass	7	13
arrowleaf groundsel	2	8
unknown	1	7
wild sasparilla	3	5
starry solomon-plume	1	4
dwarf nightshade	*	3
wild ginger	*	2
carex spp.	*	2
bracken-fern	*	2
starry solomon-plume	*	2
not present		1
pathfinder	*	1
waterleaf	*	1
mint spp.	*	1
bigroot	*	1
arrowleaf coltsfoot	*	1
alpine pyrola	*	1
false bugbane	*	1

Within eastside, forested wetland, UMAs the most frequently encountered dominant herbs were lady-fern, grass, and arrowleaf groundsel.

Table HERB-26. Eastside UMAs, upland forests, dominant herb #1 mean subplot coverage and constancy (total sites = 2, total subplots = 197).

<u>Herb Name</u>	<u>Coverage</u>	<u>Constancy</u>
pinegrass	9	22
not present		17
round-leafed violet	*	16
sidebells pyrola	*	9
bluebunch wheatgrass	3	5
northwest sedge	*	5
meadowrue	1	5
white flowered hawkweed	*	4
starry solomon-plume	*	4
aster spp.	1	3
mint spp.	*	2
broadleaf lupine	*	2
unknown	*	2
Idaho fescue	*	1
grass	*	1
western yarrow	*	1
pathfinder	*	1
lady-fern	*	1
clk sedge	*	1
beadlilly	*	1

Within eastside, upland forest, UMAs the most frequently encountered dominant herbs were pinegrass and roundleaf violet. Dominant herbs were not present in 17% of the subplots.

Table HERB-27. Eastside UMA, forested wetland, dominant herb #2 mean subplot coverage and constancy (total sites = 2, total subplots = 174).

<u>Herb Name</u>	<u>Coverage</u>	<u>Constancy</u>
lady-fern	3	20
starry solomon-plume	1	15
arrowleaf groundsel	1	12
wild sasparilla	1	6
grass	1	6
unknown	*	6
dwarf nightshade	*	6
wild ginger	*	5
sweetscented bedstraw	*	4
waterleaf	*	4
claspleaf twistedstalk	*	4
horsetail	*	2
carex spp.	*	2
mint spp.	*	2
not present		2
false bugbane	*	2
pathfinder	*	1
bluegrass spp.	*	1
alpine pyrola	*	1
pyrola spp.	*	1

Within eastside, forested wetland, UMAs the sub-dominant herbs most commonly encountered were lady-fern, starry solomon-plume, and arrowleaf groundsel.

Table HERB-28. Eastside UMA, upland forest, dominant herb #2 mean subplot coverage and constancy (total sites = 2, total subplots = 197).

<u>Herb Name</u>	<u>Coverage</u>	<u>Constancy</u>
not present		37
broadleaf lupine	1	11
round-leafed violet	*	11
pinegrass	*	5
starry solomon-plume	*	5
broadpetal strawberry	*	3
bunchgrass spp.	1	3
bigleaf sandwort	*	3
alumroot spp.	*	3
western yarrow	*	2
bunchberry dogwood	*	2
white flowered hawkweed	*	2
mint spp.	*	2
grass	*	2
unknown	*	2
heart-leaf arnica	*	1
woods strawberry	*	1
rattlesnake plantain	*	1
sidebells pyrola	*	1
meadowrue	*	1

Within eastside, upland forest, UMAs the sub-dominant herbs most commonly found were broadleaf lupine and round-leafed violet. Sub-dominant herbs were not found in 37% of the subplots.

Table HERB-29. Westside UMA, bogs, dominant herb #1 mean subplot coverage and constancy (total sites = 2, total subplots = 276).

<u>Herb Name</u>	<u>Coverage</u>	<u>Constancy</u>
false lily of the valley	4	18
beargrass	9	17
bracken-fern	7	12
not present		12
skunk cabbage	4	11
unknown	3	8
carex spp.	6	8
deer-fern	*	4
swordfern	1	4
rush spp.	1	2
trillium	*	1
western starflower	*	1

Within westside, bog, UMAs the most frequently encountered dominant herbs were false lily of the valley, beargrass, and bracken-fern.

Table HERB-30. Westside UMA, forested wetland, dominant herb #1 mean subplot coverage and constancy (total sites = 6, total subplots = 552).

<u>Herb Name</u>	<u>Coverage</u>	<u>Constancy</u>
skunk cabbage	6	14
carex spp.	6	12
swordfern	4	11
water parsely	4	9
buttercup	7	9
small fruited bulrush	6	9
lady-fern	3	8
not present		8
grass	4	6
false lily of the valley	*	4
piggyback plant	2	3
Oregon oxalis	*	2
deer-fern	*	1
Canada thistle	*	1
licorice-fern	*	1
beadlily	*	1
fireweed	*	1
soft rush	*	1
candy flower	*	1
trillium	*	1

Within westside, forested wetland, UMAs the most common dominant herbs were skunk cabbage, carex species, and swordfern.

Table HERB-31. Westside UMA, upland forest, dominant herb #1 mean subplot coverage and constancy (total sites = 11, total subplots = 886).

<u>Herb Name</u>	<u>Coverage</u>	<u>Constancy</u>
swordfern	6	21
not present		12
beargrass	2	10
grass	4	9
Oregon oxalis	2	6
piggyback plant	2	6
deer-fern	*	5
lady-fern	*	4
unknown	1	4
vanilla leaf	*	3
western starflower	*	3
bleeding heart	*	2
candy flower	*	2
Cooley's hedgenettle	*	2
dwarf nightshade	*	2
bracken-fern	*	1
inside-out-flower	*	1
false lily of the valley	*	1
sweetscented bedstraw	*	1
penstemon spp.	*	1

Within westside, upland forest, UMAs the most frequently found dominant herbs were swordfern and beargrass. Herbs were not present 12% of the time.

Table HERB-32. Westside UMA, bogs, dominant herb #2 mean subplot coverage and constancy (total sites = 6, total subplots = 273).

<u>Herb Name</u>	<u>Coverage</u>	<u>Constancy</u>
not present		39
bracken-fern	2	16
false lily of the valley	2	15
rush spp.	1	4
deer-fern	*	3
unknown	*	3
beargrass	*	3
carex spp.	*	2
skunk cabbage	*	2
trillium	*	2
bunchberry dogwood	*	2
swordfern	*	2
western starflower	*	2
sedge spp.	*	1
sundew	*	1
grass	*	1
water parsely	*	1
buttercup	*	1
lady-fern	*	1
fireweed	*	1

Within westside, bog, UMAs sub-dominant herbs were most often lacking. When sub-dominant herbs were present they were most frequently bracken-fern and false lily of the valley.

Table HERB-33. Westside UMA, forested wetlands, dominant herb #2 mean subplot coverage and constancy (total sites = 2, total subplots = 550).

<u>Herb Name</u>	<u>Coverage</u>	<u>Constancy</u>
not present		16
lady-fern	2	15
water parsely	3	12
skunk cabbage	2	9
false lily of the valley	1	8
grass	3	7
buttercup	2	5
carex spp.	1	4
swordfern	*	4
small fruited bulrush	2	3
unknown	*	3
piggyback plant	*	2
stinging nettle	*	2
Cooleye's hedgenettle	*	2
sweetscented bedstraw	*	1
leafy miterwort	*	1
licorice-fern	*	1
pioneer violet	*	1
deer-fern	*	1
bleeding heart	*	1

Within westside, forested wetland, UMAs sub-dominant herbs were most frequently lacking. When they were persent they were most commonly lady-fern and water parsley.

Table HERB-34. Westside UMA, upland forests, dominant herb #2 mean subplot coverage and constancy (total sites = 11, total subplots = 886).

<u>Herb Name</u>	<u>Coverage</u>	<u>Constancy</u>
not present		33
swordfern	*	9
lady-fern	*	7
grass	*	6
unknown	*	5
Oregon oxalis	*	4
piggyback plant	*	4
dwarf nightshade	*	3
bleeding heart	*	2
deer-fern	*	2
false lily of the valley	*	2
coolwort foam flower	*	2
vanilla leaf	*	1
inside-out-flower	*	1
bracken-fern	*	1
arrowleaf groundsel	*	1
sweetscented bedstraw	*	1
candy flower	*	1
buttercup	*	1
wood-fern	*	1

Within westside, upland forest, UMAs subdominant herbs were most frequently lacking. When they were present they were most frequently swordfern and lady-fern.

The following tables display total overstory canopy closure, total shrub coverage, total forb coverage, and total grass coverage within subplots. Site and subplot numbers are provided due to the variance of site size. Total subplot number was used to determine the mean coverages.

MEAN COVERAGE AND CONSTANCIES FOR OVERSTORY CANOPY CLOSURE, TOTAL SHRUBS, FORBS, AND GRAMINOIDS

Table UMACOVER-1. Eastside UMA Mean Coverage/Constancy for Subplot Canopy, Total Shrubs, Forbs and Graminoids. *Note: Coverage values given are in percent*

UMA TYPE	Forested Wetland	Upland Forest
Canopy	91%	77%
Shrubs	45/93	42/89
Forbs	56/99	20/80
Grass	24/74	38/48
Number of sites	1	2
Number of subplots	174	197

Table UMACOVER-2. Westside UMA Mean Coverage/Constancy for Subplot Canopy, Total Shrubs, Forbs and Graminoids. *Note: Coverage values given are in percent*

UMA TYPE	Forested Wetland	Upland Forest	Bog
Canopy	85%	90%	52%
Shrubs	52/78	56/92	83/99
Forbs	56/90	37/85	44/85
Grass	49/59	19/34	53/22
Number of sites	7	18	2
Number of subplots	579	1,465	268

Overstory subplot canopy coverage was greater in westside forested wetland UMAs than in similar eastside forested wetlands. Overstory canopy closure was higher in eastside upland forests than in westside upland forests. Shrub and grass coverages and constancies were higher within westside forested wetlands and upland forests than in similar eastside sites. Forb coverage and constancy were nearly equal between the westside and eastside forested wetland and upland UMAs.

Westside bog coverages and constancies can be found in Table UMACOVER-2.

MEAN COVERAGE AND CONSTANCY VALUES FOR WATER, ROCK, SOIL, ORGANIC GROUND COVER (OGC), DOWNED WOOD 1 (DW1), DOWNED WOOD 2 (DW2), & DOWNED WOOD 3 (DW3).

The following tables display the coverage and constancy values for total water, rock, soil, and organic ground cover. The number of subplots sampled is provided in parenthesis next to the UMA type.

Water coverage is based on open water. Rock coverage is based on exposed rock, and soil coverage is based on exposed soil. Organic ground cover includes litter, duff, mosses, lichens, and fungi. Organic ground cover does not include the downed wood coverage.

Downed wood classes are based on the amount of decay the log exhibits. Downed wood 1 logs are recently fallen trees with tight bark. Downed wood 2 logs are beginning to decay on the outside, but still have a solid center. Downed wood 3 logs are decayed throughout.

UMAs

		Eastside		Westside		
UMA Type	B	FW (174)	UF (197)	B (268)	FW (579)	UF (1465)
Water	N.A.	10/10	0/0	3/1	17/13	15/6
Rock	N.A.	0/0	15/22	21/3	6/1	26/22
Soil	N.A.	11/21	10/13	20/6	7/16	11/15
OGC	N.A.	93/100	93/100	96/99	93/99	91/99
		Eastside		Westside		
UMA Type	B	FW (174)	UF (197)	B (268)	FW (579)	UF (1465)
DW1	N.A.	10/8	7/9	9/1	9/6	11/10
DW2	N.A.	8/14	10/50	15/5	10/13	10/18
DW3	N.A.	7/12	6/21	20/31	19/36	14/29

LIVE TREE DENSITY

Tree diameter was measured in the following four inch size class intervals:

Size Class	Diameter in inches
1	0.0 - 3.9
2	4.0 - 7.9
3	8.0 - 11.9
4	12.0 - 15.9
5	16.0 - 19.9
6	20.0 - 23.9
7	24+

Data were analyzed to determine the number of trees per acre and per 1000 feet within each of their size classes. Size class analysis occurred on sizes 1-7, 2-7, 3-7, 4-7. When the last size class to be shown is 3-7 it is implied that there are no trees larger than 11.9 inches in diameter within the defined category.

To be analyzed as a live tree one of the following criteria was met: live tree - undamaged, live tree - 1/3 to 1/2 of the top broken, live tree - dead top. Minimum height was 4.5 feet. There was no minimum diameter size requirement. All trees were grouped together by size class and category.

Live tree data follows:

UMA-1. Eastside, Forested Wetland UMA Mean Tree Density - Conifers					
UMA TYPE	SIZE CLASS	TREES/ ACRE	TREES/ 1000 FT.	NUMBER OF SITES	NUMBER OF STRIPS
FW	1-7	34	112	1	11
	2-7	12	41	1	11
	3-7	5	16	1	9
	4-7	2	7	1	8

Table UMA-2. Eastside, Forested Wetland UMA Mean Tree Density - Hardwoods

UMA TYPE	SIZE CLASS	TREES/ ACRE	TREES/ 1000 FT	NUMBER OF SITES	NUMBER OF STRIPS
FW	1-7	30	100	1	12
	2-7	22	73	1	12
	3-7	19	61	1	12
	4-7	13	44	1	12

Eastside, forested wetland, UMAs had similar numbers of hardwoods and conifers per acre. Although the number of conifers are nearly equal to the number of hardwoods there were more hardwoods over size class 4.

Table UMA-3. Eastside, Upland Forest UMA Mean Tree Density - Conifers

UMA TYPE	SIZE CLASS	TREES/ ACRE	TREES/ 1000 FT	NUMBER OF SITES	NUMBER OF STRIPS
UF	1-7	96	383	2	11
	2-7	63	255	2	11
	3-7	27	108	2	11
	4-7	5	19	2	9

Table UMA-4. Eastside, Upland Forest UMA Mean Tree Density - Hardwoods

UMA TYPE	SIZE CLASS	TREES/ ACRE	TREES/ 1000 FT	NUMBER OF SITES	NUMBER OF STRIPS
UF	1-7	18	69	2	10
	2-7	16	62	2	10
	3-7	10	40	2	10
	4-7	7	25	2	7

Eastside, upland forest, UMAs had more conifers per acre than hardwoods. The conifers also were larger.

Table UMA-5. Westside, Forested Wetland UMA Mean Tree Density - Conifers

UMA TYPE	SIZE CLASS	TREES/ ACRE	TREES/ 1000 FT	NUMBER OF SITES	NUMBER OF STRIPS
FW	1-7	31	110	7	35
	2-7	14	70	7	34
	3-7	8	44	7	31
	4-7	5	24	7	27

Table UMA-6. Westside, Forested Wetland UMA Mean Tree Density - Hardwoods

UMA TYPE	SIZE CLASS	TREES/ ACRE	TREES/ 1000 FT	NUMBER OF SITES	NUMBER OF STRIPS
FW	1-7	45	135	7	36
	2-7	28	94	7	34
	3-7	11	44	7	31
	4-7	5	19	6	28

Westside, forested wetland, UMAs were dominated by hardwoods. The majority of the trees within these sites were below 12 inches in diameter.

Table UMA-7. Westside, Upland Forest UMA Mean Tree Density - Conifers

UMA TYPE	SIZE CLASS	TREES/ ACRE	TREES/ 1000 FT	NUMBER OF SITES	NUMBER OF STRIPS
UF	1-7	34	132	16	73
	2-7	18	70	16	70
	3-7	11	43	15	59
	4-7	6	26	15	51

Table UMA-8. Westside, Upland Forest UMA Mean Tree Density - Hardwoods

UMA TYPE	SIZE CLASS	TREES/ ACRE	TREES/ 1000 FT	NUMBER OF SITES	NUMBER OF STRIPS
UF	1-7	49	189	17	85
	2-7	30	130	16	83
	3-7	18	70	16	80
	4-7	9	39	15	75

Westside, upland forest, UMAs had a higher concentration of hardwoods per acre than conifers.

Table UMA-9. Westside, Bog UMA Mean Tree Density - Conifers

UMA TYPE	SIZE CLASS	TREES/ ACRE	TREES/ 1000 FT	NUMBER OF SITES	NUMBER OF STRIPS
B	1-7	29	566	2	11
	2-7	8	110	2	11
	3-7	3	22	2	11
	4-7	2	6	2	8

Table UMA-10. Westside, Bog UMA Mean Tree Density - Hardwoods

UMA TYPE	SIZE CLASS	TREES/ ACRE	TREES/ 1000 FT	NUMBER OF SITES	NUMBER OF STRIPS
B	1-7	86	335	2	12
	2-7	4	15	2	8
	3-7	1	7	1	6
	4-7	1	1	1	2

Westside, bog, UMAs contained a greater number of hardwoods per acre than conifers. Over 80% of these hardwoods were below four inches in diameter. High tree densities within bog UMAs is attributed to the ring of trees left around the actual bog post harvest.

SNAG DENSITY

Snags were defined in the following manner: recent dead (needles or leaves dead, yet still on the tree), dead tree - tight bark, or dead tree - loose bark. Minimum height was 4.5 feet. There was no minimum diameter size requirement. All snags were grouped together by size class and category.

Table UMA-11. Eastside, Forested Wetland UMA Mean Snag Density - Conifers					
UMA TYPE	SIZE CLASS	SNAGS/ ACRE	SNAGS/ 1000 FT	NUMBER OF SITES	NUMBER OF STRIPS
FW	1-7	1	5	1	6
	2-7	1	3	1	5
	3-7	1	1	1	3

Table UMA-12. Eastside, Forested Wetland UMA Mean Snag Density - Hardwoods					
UMA TYPE	SIZE CLASS	SNAGS/ ACRE	SNAGS/ 1000 FT	NUMBER OF SITES	NUMBER OF STRIPS
FW	1-7	5	17	1	8
	2-7	5	15	1	8
	3-7	3	9	1	5
	4-7	1	1	1	3

Eastside, forested wetland, UMAs contained more hardwood snags per acre than conifer snags.

Table UMA-13. Eastside, Upland Forest UMA Mean Snag Density - Conifers					
UMA TYPE	SIZE CLASS	SNAGS/ ACRE	SNAGS/ 1000 FT	NUMBER OF SITES	NUMBER OF STRIPS
UF	1-7	36	145	2	11
	2-7	24	97	2	11
	3-7	5	19	2	10
	4-7	1	2	2	4

Table UMA-14. Eastside, Upland Forest UMA Mean Snag Density - Hardwoods

UMA TYPE	SIZE CLASS	SNAGS/ ACRE	SNAGS/ 1000 FT	NUMBER OF SITES	NUMBER OF STRIPS
UF	1-7	9	33	2	9
	2-7	8	28	2	8
	3-7	2	14	1	3
	4-7	1	3	1	2

Eastside, upland forest, UMAs contained more conifer snags per acre than hardwood snags.

Table UMA-15. Westside, Forested Wetland UMA Mean Snag Density - Conifers

UMA TYPE	SIZE CLASS	SNAGS/ ACRE	SNAGS/ 1000 FT	NUMBER OF SITES	NUMBER OF STRIPS
FW	1-7	4	24	4	15
	2-7	3	19	4	14
	3-7	1	8	4	11
	4-7	1	7	3	8

Table UMA-16. Westside, Forested Wetland UMA Mean Snag Density - Hardwoods

UMA TYPE	SIZE CLASS	SNAGS/ ACRE	SNAGS/ 1000 FT	NUMBER OF SITES	NUMBER OF STRIPS
FW	1-7	5	25	6	19
	2-7	3	15	6	15
	3-7	1	6	4	8
	4-7	1	3	1	4

Westside, forested wetland, UMAs contained a similar number of hardwood snags per acre as conifer snags per acre.

Table UMA-17. Westside, Upland Forest UMA Mean Snag Density - Conifers

UMA TYPE	SIZE CLASS	SNAGS/ ACRE	SNAGS/ 1000 FT	NUMBER OF SITES	NUMBER OF STRIPS
UF	1-7	7	25	14	41
	2-7	2	11	14	35
	3-7	1	7	12	24
	4-7	1	4	10	20

Table UMA-18. Westside, Upland Forest UMA Mean Snag Density - Hardwoods

UMA TYPE	SIZE CLASS	SNAGS/ ACRE	SNAGS/ 1000 FT	NUMBER OF SITES	NUMBER OF STRIPS
UF	1-7	4	22	16	65
	2-7	3	16	16	59
	3-7	1	8	10	33
	4-7	1	4	11	22

Westside, upland forest, UMAs contained more conifer snags per acre than hardwood snags.

Table UMA-19. Westside, Bog UMA Mean Snag Density - Conifers

UMA TYPE	SIZE CLASS	SNAGS/ ACRE	SNAGS/ 1000 FT	NUMBER OF SITES	NUMBER OF STRIPS
B	1-7	1	3	2	3
	2-7	1	2	2	2

Table UMA-20. Westside, Bog UMA Mean Snag Density - Hardwoods

UMA TYPE	SIZE CLASS	SNAGS/ ACRE	SNAGS/ 1000 FT	NUMBER OF SITES	NUMBER OF STRIPS
B	1-7	4	22	16	65
	2-7	3	16	16	59
	3-7	1	8	10	33
	4-7	1	4	11	22

Westside, bog, UMAs contained more hardwood snags per acre than conifer snags.

RECOMMENDATIONS

Site Selection

To make the process of site selection more efficient, a master list of FPAs containing either RMZs/UMAS shall be requested from the DNR Forest Practice Rules and Regulations office in Olympia. Included on this list will be: FPA number, the number of UMAs, UMA acreage, water types of RMZs, length of RMZs, Township, Range, and section number, owners first and last name, and the owners phone number. Individual FPAs can then be requested from the DNR regional offices. This will eliminate the need to visit each region's office individually.

Sampling Methods

Record blowdowns in the tree data by the species, diameter at breast height, and with a "B". Record only those blowdowns that, when standing, were within the macro-plot.

Record RMZ/UMA length measured by the following formula:

$$(\# \text{ of strips sampled} \times 250 \text{ ft.}) - 250 \text{ ft.}$$

Strips are 250 ft. apart with strip # 1 beginning at zero feet, therefore the subtraction of 250 ft. Using this formula provides a more accurate representation of RMZ/UMA length sampled.

Record the distance to the nearest road in 50 foot intervals as opposed to the nearest foot.

On the east side of the state, end the sampling effort at 30 feet when the harvest boundary, due to selective cuts, is not easily identified. Where harvest unit boundary can be identified, end the sampling effort at that point.

For UMAs, record the distance to the nearest type 1, 2, 3, or 4 water in feet.

Plant Association Community Classification System

Currently Forest Service Plant Association Keys are used to characterize sampled sites. The majority of these keys were written for areas of higher elevation than we sample with little emphasis was given to riparian areas. Similar keys can be created from our data base for the lower elevation riparian zones we sampled by conducting a statistical cluster analysis to our data. These new keys could be tailored for riparian area classification. The new keys would be more accurate when applied to this project.

ACKNOWLEDGEMENTS

The following people have contributed their time to the project and deserve thanks and recognition: Chad Armour for leading the project from 1988 to 1989, Roosevelt McKenzie (WDW's Data Administrator) for his help restructuring the data base, analyzing the data and compiling the 1989 Final Report, TFW cooperators for their assistance in locating study sites, the Wildlife Steering Committee for technical advice, Rollie Geppert and John Mankowski for administrative support, and lastly but most importantly, thanks to the 1989 field data collection crew: Andy Carlson, Matt Green, Lori Braun, Amy Cook, Debbie Twigg, and Kendra Milam.

LITERATURE CITED

- Brown, Reade, E. 1985. Management of Wildlife and Fish Habitats in Forests of Western Oregon and Washington. USDA For. Serv. R6-F&WL-192-1985, Pt. 1. 323pp.
- Franklin, J. F. and C. T. Dyrness. 1973. Natural vegetation of Oregon and Washington. USDA For. Serv. Gen. Tech. Rep. PNW-8. 417pp.
- Garrison, G.A., J. M. Skovlin, C. E. Poulton, and A. H. Winward, 1976. Northwest Plant Names and Symbols for Ecosystem Inventory and Analysis. USDA For. Serv. Gen. Tech. Rep. 4th ed. 263pp.
- Henderson, J. A., D. H. Peter, R. D. Leshner, and D. C. Shaw. 1989. Forested plant associations of the Olympic National Forest. USDA For. Serv. Rep. No. R6-E-TP-001-88. 502pp.
- Leshner, R. and J. A. Henderson. 1986. A guide to the indicator species of the Olympic and Mt. Baker-Snoqualmie National Forests. USDA For. Serv. 48pp.
- Leshner, R.D. and R. H. McClure. 1986. Major indicator shrubs and herbs on national forests of Western Oregon and Southwestern Washington. USDA For. Serv. R6-TM-229-1986.
- Meuller-Dubois, D. and H. Ellenberg. 1974. Aims and methods of vegetation ecology. 547pp. John Wiley & Sons, Inc., New York.
- Thomas, J. W., C. Maser, and J. E. Rodiek. 1979. Riparian zones. In J. W. Thomas (ed.) Wildlife habitats in managed forests: The Blue Mountains of Oregon and Washington. USDA For. Serv. Agric. Handbook No. 553. pages 40-47. U. S. Gov. Printing Office
- Timber/Fish/Wildlife Agreement. 1987. A better future for our woods and streams - final report., 57pp.
- Topik, C., N. M. Halverson, and D. G. Brockway. 1986. Plant association and management guide for the western hemlock zone: Gifford Pinchot National Forest. USDA For. Serv. R6-Ecol-230B-1986. 84pp.
- United States Fish and Wildlife Service. 1986. 1986 Wetland Plant List, Washington. U.S.F.W.S. WELUT-86/W12.47. 27pp.
- Washington Department of Wildlife. 1988. Field procedures handbook - characterization of riparian management zones and upland management areas with respect to wildlife habitat. 15pp.

Washington Natural Heritage Program. 1987. State of Washington Natural Heritage Plan. WA Dept. Natural Resources. 102pp.

Washington State Forest Practices Board. 1988. Washington forest practices rules and regulations. 139pp.

Williams, C. K. and T. R. Lillybridge. 1983. Forested plant associations of the Okanogan National Forest. USDA For. Serv. Rep. No. R6-Ecol-132b-1983. 116pp.

Williams, C. K. and T. R. Lillybridge. 1985. Draft forested plant associations of the Colville National Forest. USDA For. Serv. 96pp.

Williams, C. K. and T. R. Lillybridge. 1987. Major indicator shrubs and herbs on national forests of eastern Washington. USDA For. Serv. R6-TM-TP-304-87.

APPENDIX A**LIST OF ABBREVIATIONS SCIENTIFIC AND COMMON NAMES OF TREES AND SHRUBS****TREES**

CODE	SCIENTIFIC NAME	COMMON NAME
ABAM	<i>Abies amabilis</i>	Pacific silver fir
ABGR	<i>Abies grandis</i>	grand fir
ABLA2	<i>Abies lasiocarpa</i>	subalpine fir
ABPR	<i>Abies procera</i>	noble fir
ACMA	<i>Acer macrophyllum</i>	bigleaf maple
ALRU	<i>Alnus rubra</i>	red alder
ARM	<i>Arbutus menziesii</i>	Pacific madrone
B EGL	<i>Betula occidentalis</i>	water birch
BEPA	<i>Betula papyrifera</i>	paper birch
CONU	<i>Cornus nuttallii</i>	Pacific dogwood
FRLA	<i>Fraxinus latifolia</i>	Oregon ash
LAOC	<i>Larix occidentalis</i>	Western larch
PIEN	<i>Picea engelmannii</i>	Englemann spruce
PISI	<i>Picea sitchensis</i>	Sitka spruce
PICO	<i>Pinus contorta</i>	lodgepole pine
PIMO	<i>Pinus monticola</i>	western white pine
PIPO	<i>Pinus ponderosa</i>	ponderosa pine
POTR	<i>Populus tremuloides</i>	quaking aspen
POTR2	<i>Populus trichocarpa</i>	black cottonwood
PREM	<i>Prunus emarginata</i>	bitter cherry
PSME	<i>Pseudotsuga menziesii</i>	Douglas-fir
SALIX	<i>Salix</i> spp.	willow
TABR	<i>Taxus brevifolia</i>	Pacific yew
THPL	<i>Thuja plicata</i>	western red cedar
TSHE	<i>Tsuga heterophylla</i>	western hemlock
TSME	<i>Tsuga mertensiana</i>	mountain hemlock

SHRUBS

CODE	SCIENTIFIC NAME	COMMON NAME
ACCI	<i>Acer circinatum</i>	vine maple
ACGLD	<i>Acer glabrum</i>	Douglas maple var. douglasii
ALIN	<i>Alnus incana</i>	mountain alder
ALNUS	<i>Alnus</i> spp.	alder
ALSI	<i>Alnus sinuata</i>	Sitka alder
AMAL	<i>Amalanchier alnifolia</i>	serviceberry
ARUV	<i>Arctostaphylos uva-ursi</i>	bearberry
BENE	<i>Berberis nervosa</i>	Cascade Ore grape
BERE	<i>Berberis repens</i>	Oregon grape
CEVE	<i>Ceanothus velutinus</i>	snowbrush ceanothus
CHUM	<i>Chimaphyla umbellata</i>	common prince's-pine
CONU	<i>Cornus nuttallii</i>	pacific dogwood
COST	<i>Cornus stolonifera</i>	red-osier dogwood
COCO2	<i>Corylus cornuta</i>	hazelnut
CRDO	<i>Crataegus douglasii</i>	black hawthorn
CYSC	<i>Cytisus scoparius</i>	Scot's broom
GASH	<i>Gaultheria shallon</i>	salal
HODI	<i>Holodiscus discolor</i>	ocean-spray
HOLLY	<i>Ilex</i> spp.	holly
KAOC	<i>Kalmia occidentalis</i>	swamp laurel
LEGL	<i>Ledum glandulosum</i>	smooth Labrador-tea
LIBOL	<i>Linnaea borealis</i>	twinflower var. longiflora
LONIC	<i>Lonicera</i> spp.	honeysuckle
LOIN	<i>Lonicera involucrata</i>	black twin-berry
LOUT2	<i>Lonicera utahensis</i>	Utah honeysuckle
MEFE	<i>Menziesia ferruginea</i>	rusty menziesia
OECE	<i>Oemleria cerasiformis</i>	Indian plum
OPHO	<i>Oplopanax horridum</i>	devil's club
PAMY	<i>Pachistima myrsinities</i>	pachistima
PHLE2	<i>Philadelphus lewisii</i>	mockorange
PHCA3	<i>Physocarpus capitatus</i>	Pacific ninebark
PHMA	<i>Physocarpus malvaceus</i>	mallow ninebark
PREM	<i>Prunus emarginata</i>	bittercherry
PRVI	<i>Prunus virginiana</i>	common chokecherry
PYFU	<i>Pyrus fusca</i>	western crabapple
RHPU	<i>Rhamnus purshiana</i>	cascara
RHAL	<i>Rhododendron albiflorum</i>	white rhododendron
RHDI	<i>Rhus diversiloba</i>	poison-ivy
RIBES	<i>Ribes</i> spp.	currant

RIBR	<i>Ribes bracteosum</i>	stink currant
RICE	<i>Ribes cereum</i>	wax currant
RILA	<i>Ribes lacustre</i>	prickly currant
RIVI	<i>Ribes viscosissimum</i>	sticky currant
ROSA	<i>Rosa</i> spp.	rose
ROGY	<i>Rosa gymnocarpa</i>	baldhip rose
RONUH	<i>Rosa nutcana</i> var. <i>hispida</i>	bristly Nootka rose
ROWO	<i>Rosa woodsii</i>	Wood's rose
RUBUS	<i>Rubus</i> spp.	rubus
RUDI	<i>Rubus discolor</i>	Himalayan blackberry
RULA	<i>Rubus lasiococcus</i>	dwarf bramble
RULE	<i>Rubus leucodermis</i>	blackcap
RUPA	<i>Rubus parviflorus</i>	westrn thimbleberry
RUSP	<i>Rubus spectabilis</i>	salmonberry
RUURU	<i>Rubus ursinus</i>	trailing blackberry
SALIX	<i>Salix</i> spp.	Willow
SASC	<i>Salix scouleriana</i>	Scouler willow
SACE	<i>Sambucus cerulea</i>	blue elderberry
SARA	<i>Sambucus racemosa</i>	red elderberry
SHCA	<i>Shepherdia canadensis</i>	russet buffaloberry
SOSC2	<i>Sorbus scopulina</i>	mountain ash
SPBEL	<i>Spirea betulifolia</i>	shiny leaf spirea var. lucinda
SPDO	<i>Spirea douglasii</i>	hardhack
SYAL	<i>Symphoricarpos albus</i>	common snowberry
SYMOH	<i>Symphoricarpos mollis</i>	creeping snowberry var. hesperius
VACCI	<i>Vaccinium</i> spp.	huckleberry
VAAL	<i>Vaccinium alaskaense</i>	Alaska huckleberry
VAME	<i>Vaccinium membranaceum</i>	big huckleberry
VAMY	<i>Vaccinium myrtillus</i>	low huckleberry
VAOV2	<i>Vaccinium ovatum</i>	evergreen huckleberry
VAPA	<i>Vaccinium parvifolium</i>	red huckleberry
VASC	<i>Vaccinium scoparium</i>	grouse huckleberry

HERBS

CODE	SCIENTIFIC NAME	COMMON NAME
ACMI	<i>Achillea millefolium</i>	common yarrow
ACRU	<i>Actaea rubra</i>	baneberry
ACTR	<i>Achlys triphylla</i>	vanilla leaf
ADBI	<i>Adenocaulon bicolor</i>	pathfinder
ADPE	<i>Adiantum pedatum</i>	maidenhair fern
AGUR	<i>Agastache urticifolia</i>	nettle-leaf horse-mint
AGSP	<i>Agropyron spicatum</i>	bluebunch wheatgrass
ANMA	<i>Anaphalis margaritacea</i>	pearly-everlasting
ARNU3	<i>Aralia nudicaulis</i>	wild sasparilla
ARMA3	<i>Arenaria macrophylla</i>	bigleaf sandwort
ARCO	<i>Arnica cordifolia</i>	heart-leaf arnica
ARSY	<i>Aruncus sylvester</i>	goatsbeard
ASCA3	<i>Asarum caudatum</i>	wild ginger
ASTER	<i>Aster</i> spp.	Aster
ASCO	<i>Aster conspicuus</i>	showy aster
ATFI	<i>Athyrium filix-femina</i>	lady-fern
BLSP	<i>Blechnum spicant</i>	deerfern
BROMU	<i>Bromus</i> spp.	brome
BRBR	<i>Bromus brizaeformis</i>	rattle grass
BRVU	<i>Bromus vulgaris</i>	Columbia brome
CARO	<i>Campanula rotundifolia</i>	harebell
CARU	<i>Calamagrostis rubescens</i>	pinegrass
CAREX	<i>Carex</i> spp.	carex
CACO	<i>Carex concinnoides</i>	northwest sedge
CAGE	<i>Carex geeyeri</i>	clk sedge
CASTI	<i>Castilleja</i> spp.	Indian-paintbrush
CIAR	<i>Cirsium arvense</i>	Canada thistle
CIRSI	<i>Cirsium</i> spp.	thistle
CIVU	<i>Cirsium vulgare</i>	bull thistle
CLCOL	<i>Clematis columbiana</i>	Columbia clematis
CLUN	<i>Clintonia uniflora</i>	beadlily
COCA	<i>Cornus canadensis</i>	bunchberry dogwood
COSC	<i>Corydalis scouleri</i>	Scouler's corydalis
CYMO	<i>Cypripedium montanum</i>	mountain lady's-slipper
DAGL	<i>Dactylis glomerata</i>	orchard-grass
DELPH	<i>Delphinium</i> spp.	larkspur
DIFO	<i>Dicentra formosa</i>	bleeding heart
DIPU	<i>Digitalis purpurea</i>	foxglove
DIHO	<i>Disporum hookeri</i>	Hooker fairy-bell

DITR	<i>Disporum trachycarpum</i>	wartberry fairy-bell
DRRO	<i>Drosera rotundifolia</i>	sundew
DRAU2	<i>Dryopteris austriaca</i>	wood-fern
EPAN	<i>Epilobium angustifolium</i>	fireweed
ERIGE	<i>Erigeron</i> spp.	daisy
EQUIS	<i>Equisetum</i> spp.	horsetail
EQAR	<i>Equisetum arvense</i>	common horsetail
FEID	<i>Festuca idahoensis</i>	Idaho fescue
FRAGA	<i>Fragaria</i> spp.	strawberry
FRVE	<i>Fragaria vesca</i>	woods strawberry
FRVI	<i>Fragaria virginiana</i>	broadpetal strawberry
GABO	<i>Galium boreale</i>	northern bedstraw
GATR	<i>Galium triflorum</i>	sweetscented bedstraw
GEVI	<i>Geranium viscosissimum</i>	sticky purple geranium
GEUM	<i>Geum macrophyllum</i>	Oregon avens
GLHE	<i>Glecoma hederacea</i>	ground ivy
GOOB	<i>Goodyera oblongifolia</i>	western rattlesnake plain-tain
GYDR	<i>Gymnocarpium dryopteris</i>	oakfern
HELA	<i>Heracleum lanatum</i>	cow-parsnip
HEMI	<i>Heuchera micrantha</i>	alumroot
HEUCH	<i>Heuchera</i> spp.	alumroot
HIAL	<i>Hieracium albiflorum</i>	white-flowered hawkweed
HYTE	<i>Hydrophyllum tenuipes</i>	waterleaf
JUNUC	<i>Juncus</i> spp.	rush
JUEFE	<i>Juncus effusus</i>	soft rush
LACTU	<i>Lactuca</i> spp.	lettuce
LAMU	<i>Lactuca muralis</i>	wall lettuce
LOMAT	<i>Lomatium</i> spp.	biscuit-root
LUPIN	<i>Lupinus</i> spp.	lupine
LULAS	<i>Lupinus latifolius</i>	broadleaf lupine
LUSEA	<i>Lupinus sericeus</i>	silky lupine
LUZSP	<i>Luzula</i> spp.	woodrush
LYCL	<i>Lycopodium clavatum</i>	stag's horn moss
LYAM	<i>Lysichitum americanum</i>	skunk cabbage
MADI2	<i>Maianthemum dilatatum</i>	false lilly of the valley
MAOR	<i>Marah oreganus</i>	bigroot
MELI	<i>Melampyrum lineare</i>	cow-wheat
MECI	<i>Mentha citrata</i>	bergamot mint
MENTH	<i>Mentha</i> spp.	mint
MILE	<i>Mimulus lewisii</i>	Lewis' monkey-flower
MIGU	<i>Mimulus guttatus</i>	yellow monkey-flower

MICA3	<i>Mitella caulescens</i>	leafy mitrewort
MOSI	<i>Montia sibirica</i>	miner's lettuce
OESA	<i>Oenanthe sarmentosa</i>	water-parsley
OSCH	<i>Osmorhiza chilensis</i>	mountain sweet-root
OXOR	<i>Oxalis oregana</i>	Oregon oxalis
PEBRA	<i>Pedicularis bracteosa</i>	bracted lousewort
PEFR	<i>Petasites frigidus</i>	coltsfoot
PENST	<i>Penstemon</i> spp.	beardtongue
PESA	<i>Petasites sagittatus</i>	arrowleaf coltsfoot
PHAR	<i>Phalaris arundinacea</i>	canarygrass
PLRE	<i>Pleuropogon refractus</i>	nodding semaphoregrass
POGL	<i>Polypodium glycyrrhiza</i>	licorice-fern
POMU	<i>Polystichum munitum</i>	swordfern
POA	<i>Poa</i> spp.	bluegrass
PRVU	<i>Prunella vulgaris</i>	self-heal
PTAQ	<i>Pteridium aquilinum</i>	bracken fern
PYAS	<i>Pyrola asarifolia</i>	alpine pyrola
PYPI	<i>Pyrola picta</i>	white vein pyrola
PYSE	<i>Pyrola secunda</i>	sidebells pyrola
RANUN	<i>Ranunculus</i> spp.	buttercup
RUMEX	<i>Rumex</i> spp.	dock
SAAC	<i>Satureja acinos</i>	savory
SCMI	<i>Scirpus microcarpus</i>	small-fruited bulrush
SESP	<i>Sedum spathulifolium</i>	broadleaf stonecrop
SEJA	<i>Senecio jacobaea</i>	tansy ragwort
SETR	<i>Senecio triangularis</i>	arrowleaf groundsel
SODU2	<i>Solanum dulcamara</i>	climbing nightshade
SOCA	<i>Solidago canadensis</i>	meadow goldenrod
SMRA	<i>Smilacina racemosa</i>	western Solomon-plume
SMST	<i>Smilacina stellata</i>	starry solomon-plume
STCO4	<i>Stachys cooleyae</i>	Cooley's betony
STAM	<i>Streptopus amplexifolius</i>	claspleaf twistedstalk
STRO	<i>Streptopus rosea</i>	rosey twisted-stalk
TARAX	<i>Taraxacum</i> spp.	dandelion
TAOF	<i>Taraxacum officinale</i>	common dandelion
TEGR	<i>Tellima grandiflora</i>	fringecup
THOC	<i>Thalictrum occidentale</i>	meadowrue
TITR	<i>Tiarella trifoliata</i>	coolwort foamflower
TOME	<i>Tolmiea menziesii</i>	piggyback plant
TRCA3	<i>Trautvetteria caroliniensis</i>	false bugbane
TRLA2	<i>Trientalis latifolia</i>	western starflower
TRIFO	<i>Trifolium</i> spp.	clover

TROV	<i>Trillium ovatum</i>	trillium
TYLA	<i>Typha latifolia</i>	common cat-tail
VAHE	<i>Vancouveria hexandra</i>	inside-out-flower
VECA	<i>Veratrum californicum</i>	California false hellebore
VIOLA	<i>Viola</i> spp.	violet
VIGL	<i>Viola glabrella</i>	pioneer violet
VIOR2	<i>Viola orbiculata</i>	round-leaved violet
URDI	<i>Urtica dioica</i>	stinging nettle
XETE	<i>Xerophyllum tenax</i>	beargrass

APPENDIX B**KEY CONTACTS: SOURCE FOR FOREST PRACTICE INFORMATION****DEPARTMENT OF NATURAL RESOURCES**

<u>REGION</u>	<u>NAME</u>	<u>TITLE</u>	<u>TELEPHONE</u>
CEN	John Baarspul	FP Regional Coordinator	(206) 753-3410
CEN	Debie Boyd	FP Admin Asst	(206) 753-3410
NE	Bob Anderson	FP Regional Coordinator	(509) 684-5201
NE	Bob Hartley	Deer Park FP Forester	(509) 684-5201
NE	Al Lang	Chewelah FP Forester	(509) 684-5201
NE	Diana Hoffman	FP Admin Asst	(509) 684-5201
NE	Mel Kuipers	Republic FP Forester	(509) 684-5201
NE	Don Strand	Colville FP Forester	(509) 684-5201
NW	Dave Dietzman	FP Regional Coordinator	(206) 856-0083
NW	Diane Paustain	FP Admin Asst	(206) 856-0083
OLY	Russ Holt	Sequim FP Forester	(206) 374-6131
OLY	Dan Christensen	Ozette FP Forester	(206) 374-6131
OLY	Wayne Radcliff	Quinalt FP Forester	(206) 288-2448
OLY	Jackie Simmons	FP Admin Asst	(206) 374-6131
OLY	Jack Zaccardo	FP Regional Coordinator	(206) 374-6131
SPS	Diane Andersen	FP Admin Asst	(206) 825-1631
SPS	Ben Cleveland	FP Regional Coordinator	(206) 825-1631
SE	Don Aden	South Half FP Forester	(509) 962-1006
SE	Linda Hazlett	FP Admin Asst	(509) 925-6131
SE	Len Rigglin	North Half FP Forester	(509) 962-1006
SE	Ben Startt	FP Regional Coordinator	(509) 925-6131
SW	Llyod Handlos	FP Regional Coordinator	(206) 577-2025
SW	Shirley Shea	FP Admin Asst	(206) 577-2025

WEYERHAEUSER

<u>REGION</u>	<u>NAME</u>	<u>TITLE</u>	<u>TELEPHONE</u>
CEN	John Helm	Area Forester	(206) 748-8661
CEN	Ken Lentz	District Engineer	(206) 748-1167
CEN	Kieth Metcalf	District Engineer	(206) 942-2442
CEN	Tim Shere	District Engineer	(206) 942-2442
CEN	Warren Sorenson	District Engineer	(206) 748-8661
OLY	Don Jordan	District Engineer	(206) 532-7110
SPS	Steve Anderson	TFW Industry Coord.	(206) 888-2511
SPS	Mike Bradley	Area Forester	(206) 825-5715
SW	John Keatly	TFW Industry Coord.	(206) 425-2150

SW	Jim Booher	District Engineer	(206) 425-2150
----	------------	-------------------	----------------

PLUM CREEK

<u>REGION</u>	<u>NAME</u>	<u>TITLE</u>	<u>TELEPHONE</u>
NE	Dwight Opp	Timberlands Superint.	(509) 447-3686
SPS	Gary Johnson	Timberlands Superint.	(206) 825-5596
SE	Pete Heide	Timberlands Superint.	(509) 649-2218
SE	Steve Griswold	Forester	(509) 649-2218
SW	Roger Wimer	Production Superint.	(206) 636-2650

OTHER INDUSTRY

<u>REGION</u>	<u>NAME</u>	<u>COMPANY</u>	<u>TELEPHONE</u>
CEN	Al Cain	Campbell Group	(206) 532-7331
CEN	John Ensinger	Menesha	(206) 754-1711
CEN	Bob Schwarz	Murray Pacific	(206) 492-5981
NE	Steve Tveit	Boise Cascade	(509) 738-6421
NE	Wayne Vaagen	Vaagen Bros.	(509) 684-5071
NW	Dave Chaimberlain	Georgia Pacific	(206) 733-4410
NW	Pete Poeschol	Poeschol & Schultz	(206) 659-5666
NW	Bill Rawlins	Crown Pacific	(206) 826-3951
NW	Norm Schaaf	Crown Pacific	(206) 826-3951
OLY	Frank Phillips	ITT Rayonier	(206) 374-6565
SPS	Craig Beals	Champion International	(206) 879-5311
SPS	Vaughn Webb	Pope Resources	(206) 297-3341
SPS	Mike Masman	PBMCO Land Trust	(206) 624-5810
SPS	Dave Baxtrum	Simpson Timber	(206) 426-3381
SE	Jeff Davies	Boise Cascade	(206) 925-5341
SE	Bill Hatch	Boise Cascade	(509) 773-4343
SE	Bill Howard	Boise Cascade	(509) 453-3131
SE	Jeff Jones	Boise Cascade	(509) 925-5341
SE	Bob McGruder	Boise Cascade	(509) 925-5341
SW	Marc Norberg	International Paper	(206) 423-2110
SW	Monte Martinsen	Longview Fibre	(206) 425-1550

DEPARTMENT OF WILDLIFE

<u>REGION</u>	<u>NAME</u>	<u>TITLE</u>	<u>TELEPHONE</u>
I	John Whalen	TFW Biologist	(509) 456-4082
II	John Rohrer	TFW Biologist	(509) 754-4624
III	Bill Weiler	TFW Biologist	(509) 575-2740
IV	Dana Base	TFW Biologist	(509) 629-2488
V	Bob Bicknell	TFW Biologist	(206) 274-9814
VI	Gloria Mitchell	TFW Biologist	(206) 753-2600

October 1990

HQ	Andy Carlson	TFW Biologist	(206) 753-3318
HQ	John Mankowski	TFW Program Manager	(206) 753-3318
HQ	Pete Haug	Systems Biologist	(206) 753-3318

DEPARTMENT OF REVENUE

<u>NAME</u>	<u>TITLE</u>	<u>TELEPHONE</u>
Joyce Fouts	Systems Analyst	(206) 753-5573